

- (i) The name of the additive.
 - (ii) A statement of the concentration or the strength of the additive in any intermediate premixes.
- (2) The label or labeling of the food additive shall also bear adequate directions for use.

§ 172.808 Copolymer condensates of ethylene oxide and propylene oxide.

Copolymer condensates of ethylene oxide and propylene oxide may be safely used in food under the following prescribed conditions:

(a) The additive consists of one of the following:

(1) α -Hydro-*omega*-hydroxy-poly (oxyethylene) poly(oxypropylene)-(55-61 moles)poly(oxyethylene) block copolymer, having a molecular weight range of 9,760-13,200 and a cloud point above 100 °C in 1 percent aqueous solution.

(2) α -Hydro-*omega*-hydroxy-poly (oxyethylene)poly(oxypropylene)-(53-59 moles)poly(oxyethylene)(14-16 moles) block copolymer, having a molecular weight range of 3,500-4,125 and a cloud point of 9 °C-12 °C in 10 percent aqueous solution.

(3) α -Hydro-*omega*-hydroxy-poly(oxyethylene)/poly(oxypropylene) (minimum 15 moles)/poly(oxyethylene) block copolymer, having a minimum average molecular weight of 1900 and a minimum cloud point of 9 °C-12 °C in 10 percent aqueous solution.

(4) α -Hydro-*omega*-hydroxy-poly(oxyethylene) poly (oxypropylene)-(51-57 moles) poly(oxyethylene) block copolymer, having an average molecular weight of 14,000 and a cloud point above 100 °C in 1 percent aqueous solution.

(b) The additive is used or intended for use as follows:

(1) The additive identified in paragraph (a)(1) of this section is used in practice as a solubilizing and stabilizing agent in flavor concentrates (containing authorized flavoring oils) for use in foods for which standards of identity established under section 401 of the Act do not preclude such use, provided that the weight of the additive does not exceed the weight of the flavoring oils in the flavor concentrate.

(2) The additive identified in paragraph (a)(2) of this section is used as a processing aid and wetting agent in combination with dioctyl sodium sulfo-

succinate for fumaric acid as prescribed in §172.810.

(3) The additive identified in paragraph (a)(3) of this section is used:

(i) As a surfactant and defoaming agent, at levels not to exceed 0.05 percent by weight, in scald baths for poultry defeathering, followed by potable water rinse. The temperatures of the scald baths shall be not less than 125 °F.

(ii) As a foam control and rinse adjunct in hog dehairing machines at a use level of not more than 5 grams per hog.

(4) The additive identified in paragraph (a)(4) of this section is used as a dough conditioner in yeast-leavened bakery products for which standards of identity established under section 401 of the Act do not preclude such use, provided that the amount of the additive dose not exceed 0.5 percent by weight of the flour used.

[42 FR 14491, Mar. 15, 1977, as amended at 46 FR 57476, Nov. 24, 1981]

§ 172.809 Curdlan.

Curdlan may be safely used in accordance with the following conditions:

(a) Curdlan is a high molecular weight polymer of glucose (β -1,3-glucan; CAS Reg. No. 54724-00-4) produced by pure culture fermentation from the nonpathogenic and nontoxicogenic bacterium *Alcaligenes faecalis* var. *myxogenes*.

(b) Curdlan meets the following specifications when it is tested according to the methods described or referenced in the document entitled "Analytical Methods for Specification Tests for Curdlan," by Takeda Chemical Industries, Ltd., 12-10 Nihonbashi, 2-Chome, Chuo-ku, Tokyo, 103, Japan, 1996, which is incorporated by reference in accordance with 5 U.S.C. 552(a) and 1 CFR part 51. Copies are available from the Division of Petition Control (HFS-215), Center for Food Safety and Applied Nutrition, Food and Drug Administration, 200 C St. SW., Washington, DC 20204, or may be examined at the Center for Food Safety and Applied Nutrition's Library, Food and Drug Administration, 200 C St. SW., rm. 3321, Washington, DC, or at the Office of the Federal Register, 800 North Capitol St. NW., suite 700, Washington, DC.

§ 172.810

21 CFR Ch. I (4-1-01 Edition)

- (1) Positive for curdlan.
- (2) Assay for curdlan (calculated as anhydrous glucose), not less than 80 percent.
- (3) pH of 1 percent aqueous suspension, 6.0-7.5.
- (4) Lead, not more than 0.5 mg/kg.
- (5) Heavy metals (as Pb), not more than 0.002 percent.
- (6) Total nitrogen, not more than 0.2 percent.
- (7) Loss on drying, not more than 10 percent.
- (8) Residue on ignition, not more than 6 percent.
- (9) Gel strength of 2 percent aqueous suspension, not less than 600×10^3 dyne per square centimeter.
- (10) Aerobic plate count, not more than 10^3 per gram.
- (11) Coliform bacteria, not more than 3 per gram.

(c) Curdlan is used or intended for use in accordance with good manufacturing practice as a formulation aid, processing aid, stabilizer and thickener, and texturizer in foods for which standards of identity established under section 401 of the act do not preclude such use.

[61 FR 65941, Dec. 16, 1996]

§ 172.810 Dioctyl sodium sulfosuccinate.

The food additive dioctyl sodium sulfosuccinate, which meets the specifications of the Food Chemicals Codex, 3d Ed. (1981), pp. 102-104, which is incorporated by reference (copies may be obtained from the National Academy Press, 2101 Constitution Ave. NW., Washington, DC 20418, or may be examined at the Office of the Federal Register, 800 North Capitol Street, NW., suite 700, Washington, DC 20408), may be safely used in food in accordance with the following prescribed conditions:

(a) As a wetting agent in the following fumaric acid-acidulated foods: Dry gelatin dessert, dry beverage base, and fruit juice drinks, when standards of identity do not preclude such use. The labeling of the dry gelatin dessert and dry beverage base shall bear adequate directions for use, and the additive shall be used in such an amount that the finished gelatin dessert will contain not in excess of 15 parts per

million of the additive and the finished beverage or fruit juice drink will contain not in excess of 10 parts per million of the additive.

(b) As a processing aid in sugar factories in the production of unrefined cane sugar, in an amount not in excess of 0.5 part per million of the additive per percentage point of sucrose in the juice, syrup, or massecuite being processed, and so used that the final molasses will contain no more than 25 parts per million of the additive.

(c) As a solubilizing agent on gums and hydrophilic colloids to be used in food as stabilizing and thickening agents, when standards of identity do not preclude such use. The additive is used in an amount not to exceed 0.5 percent by weight of the gums or hydrophilic colloids.

(d) As an emulsifying agent for cocoa fat in noncarbonated beverages containing cocoa, whereby the amount of the additive does not exceed 25 parts per million of the finished beverage.

(e) As a dispersing agent in "cocoa with dioctyl sodium sulfosuccinate for manufacturing" that conforms to the provisions of §163.117 of this chapter and the use limitations prescribed in §172.520, in an amount not to exceed 0.4 percent by weight thereof.

(f) As a processing aid and wetting agent in combination with α -hydroxy- ω -hydroxy - poly(oxyethylene) - poly-(oxypropylene) (53-59 moles) poly(oxyethylene) (14-16 moles) block copolymer, having a molecular weight range of 3,500-4,125 and a cloud point of 9 °C-12 °C in 10 percent aqueous solution, for fumaric acid used in fumaric acid-acidulated dry beverage base and in fumaric acid-acidulated fruit juice drinks, when standards of identity do not preclude such use. The labeling of the dry beverage base shall bear adequate directions for use, and the additives shall be used in such an amount that the finished beverage or fruit juice drink will contain not in excess of a total of 10 parts per million of the dioctyl sodium sulfosuccinate-block copolymer combination.

[42 FR 14491, Mar. 15, 1977, as amended at 49 FR 10105, Mar. 19, 1984]