

§ 172.833

the use of the additive which, if followed, will result in a food that complies with the requirements of this section.

§ 172.833 Sucrose acetate isobutyrate (SAIB).

Sucrose acetate isobutyrate may be safely used in foods in accordance with the following prescribed conditions:

(a) Sucrose acetate isobutyrate (CAS Reg. No. 27216-37-1), or SAIB, is the chemical *alpha*-D-glucopyranoside, O-acetyl-tris-O-(2-methyl-1-oxopropyl)-*beta*-D-fructofuranosyl, acetate tris(2-methyl propanoate).

(b) SAIB, a pale, straw-colored liquid, meets the following specifications: (1) Assay: Not less than 98.8 percent and not more than 101.9 percent, based on the following formula:

$$\text{Assay} = ((\text{SV} \cdot 0.10586) + 56.1) \times 100$$

Where SV = Saponification value

(2) Saponification value: 524-540 determined using 1 gram of sample by the "Guide to Specifications for General Notices, General Analytical Techniques, Identification Tests, Test Solutions, and Other Reference Materials," in the "Compendium of Food Additive Specifications, Addendum 4, Food and Agriculture Organization of the United Nations (FAO), Food and Nutrition Paper 5, Revision 2" (1991), pp. 203 and 204, which is incorporated by reference, in accordance with 5 U.S.C. 552(a) and 1 CFR part 51. Copies are available from the Office of Premarket Approval, Center for Food Safety and Applied Nutrition (HFS-200), Food and Drug Administration, 200 C St. SW., Washington, DC 20204, or may be examined at the Center for Food Safety and Applied Nutrition's Library, 200 C St. SW., rm. 3321, Washington, DC, or at the Office of the Federal Register, 800 North Capitol St. NW., suite 700, Washington, DC.

(3) Acid value: Not to exceed 0.20 determined using 50 grams of sample by the "Guide to Specifications for General Notices, General Analytical Techniques, Identification Tests, Test Solutions, and Other Reference Materials," in the "Compendium of Food Additive Specifications, Addendum 4, FAO Food and Nutrition Paper 5, Revision 2," p. 189 (1991), which is incorporated by reference; see paragraph (b)(2) of this section for availability of the incorporation by reference.

21 CFR Ch. I (4-1-01 Edition)

tion for availability of the incorporation by reference.

(4) Lead: Not to exceed 1.0 milligrams/kilogram determined by the "Atomic Absorption Spectrophotometric Graphite Furnace Method, Method I," in the "Food Chemicals Codex," 4th ed. (1996), pp. 763 and 764, with an attached modification to the sample digestion section in Appendix III.B (July 1996), which is incorporated by reference. Copies are available from the National Academy Press, 2101 Constitution Ave. NW., Box 285, Washington, DC 20055 (Internet "http://www.nap.edu"), or may be examined at the Center for Food Safety and Applied Nutrition's Library, 200 C St. SW., rm. 3321, Washington, DC, or at the Office of the Federal Register, 800 North Capitol St. NW., suite 700, Washington, DC.

(5) Triacetin: Not to exceed 0.10 percent determined by gas chromatography as described in the "Guide to Specifications for General Notices, General Analytical Techniques, Identification Tests, Test Solutions, and Other Reference Materials," in the "Compendium of Food Additive Specifications, Addendum 4, FAO Food and Nutrition Paper 5, Revision 2," (1991), pp. 13-26, which is incorporated by reference; see paragraph (b)(2) of this section for availability of the incorporation by reference.

(c) The food additive is used as a stabilizer (as defined in § 170.3(o)(28) of this chapter) of emulsions of flavoring oils in nonalcoholic beverages.

(d) The total SAIB content of a beverage containing the additive does not exceed 300 milligrams/kilogram of the finished beverage.

[64 FR 29958, June 4, 1999; 64 FR 43072, Aug. 9, 1999]

§ 172.834 Ethoxylated mono- and diglycerides.

The food additive ethoxylated mono- and diglycerides (polyoxyethylene (20) mono- and diglycerides of fatty acids) (polyglycerate 60) may be safely used in food in accordance with the following prescribed conditions:

(a) The food additive is manufactured by:

(1) Glycerolysis of edible fats primarily composed of stearic, palmitic, and myristic acids; or