

§ 520.1205 Kanamycin sulfate, pectin, bismuth subcarbonate, activated attapulgitic tablets.

(a) *Specifications.* Each tablet contains 100 milligrams of kanamycin (as the sulfate), 25 milligrams of pectin, 250 milligrams of bismuth subcarbonate, and 500 milligrams of activated attapulgitic.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Amount.* One tablet per 44 kilograms (20 pounds) of body weight every 8 hours. Maximum dose 3 tablets every 8 hours. For animals under 22 kilograms (10 pounds) $\frac{1}{2}$ tablet every 8 hours. The initial loading dose should be twice the amount of a single dose.

(2) *Indications for use.* For the treatment of bacterial enteritis caused by organisms susceptible to kanamycin and the symptomatic relief of associated diarrhea in dogs.

(3) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[56 FR 8710, Mar. 1, 1991]

§ 520.1242 Levamisole hydrochloride oral dosage forms.

§ 520.1242a Levamisole hydrochloride drench and drinking water.

(a) *Specifications.* Each package contains either 9.075, 11.7, 18.15, 46.8, or 544.5 grams of levamisole hydrochloride.

(b) *Sponsors.* Approval for sponsors in 21 CFR 510.600(c) for use as in paragraph (d) of this section as follows:

(1) See 043781 for use of 46.8 gram package as in paragraph (d)(1) of this section, for 11.7 and 46.8 gram packages as in paragraph (d)(2) of this section, and for 9.075 and 18.15 gram packages as in paragraph (d)(3) of this section.

(2) See 000061 for use of 46.8 and 544.5 gram packages as in paragraph (d)(1) of this section, for 11.7, 46.8, and 544.5 gram packages as in paragraph (d)(2) of this section, and for 18.15 gram package as in paragraph (d)(3) of this section.

(3) See 057561 for use of 46.8 and 544.5 gram packages as in paragraphs (d)(1) and (d)(2) of this section.

(c) *Related tolerances.* See § 556.350 of this chapter.

(d) *Conditions of use.* It is used as an anthelmintic at 0.365 gram per 100 pounds of body weight as follows:

(1) *Cattle—(i) Amount.* As a single oral dose drench using 46.8 or 544.5 gram packet.

(ii) *Indications for use.* Anthelmintic effective against the following nematode infections: Stomach worms (*Haemonchus*, *Trichostrongylus*, *Ostertagia*), intestinal worms (*Trichostrongylus*, *Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*), and lungworms (*Dictyocaulus*).

(iii) *Limitations.* Conditions of constant helminth exposure may require retreatment within 2 to 4 weeks after the first treatment. Do not slaughter for food within 48 hours of treatment. Not for use in dairy animals of breeding age. Consult your veterinarian before using in severely debilitated animals. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. Prepare solutions for use as follows:

(a) Dissolve contents of 46.8 gram package in water to provide 1 quart (32 fluid ounces) of drench solution and administer as a drench at $\frac{1}{4}$ ounce per 100 pounds of body weight as a single oral dose.

(b) Dissolve contents of 46.8 gram package in water to provide 8.75 fluid ounces of concentrate solution and administer as a drench at 2 milliliters per 100 pounds of body weight as a single oral dose by syringe.

(c) Dissolve contents of 544.5 gram package in 3 liters of water and administer as a drench at 2 milliliters per 100 pounds of body weight as a single oral dose.

(2) *Sheep—(i) Amount.* As a single oral dose drench using 11.7, 46.8, or 544.5 gram packet.

(a) *Indications for use.* Anthelmintic effective against the following nematode infections: Stomach worms (*Haemonchus*, *Trichostrongylus*, *Ostertagia*), intestinal worms (*Trichostrongylus*, *Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*, *Chabertia*), and lungworms (*Dictyocaulus*).

(b) *Limitations.* Dissolve in 1 gallon (128 fluid ounces) of water and administer as a single drench at 1 ounce (0.365