

veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(2) *Turkeys*—(i) *Amount*. 100 milligrams per bird up to 12 weeks and 200 milligrams per bird over 12 weeks.

(ii) *Indications for use*. For removal of large roundworm (*Ascaridia* spp.).

(iii) *Limitations*. For use in drinking water or feed. Use as sole source of drinking water. Prepare fresh solution daily. Use as 1-day single treatment. Withdraw 14 days prior to slaughter. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(3) *Swine*—(i) *Amount*. 50 milligrams per pound of body weight.

(ii) *Indications for use*. For removal of large roundworm (*Ascaris suum*) and nodular worms (*Oesophagostomum* spp.).

(iii) *Limitations*. For use in drinking water or feed. Use as sole source of drinking water. Prepare fresh solution daily. Use as 1-day single treatment. Withdraw 21 days prior to slaughter. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[64 FR 23018, Apr. 29, 1999]

§ 520.1840 Poloxalene.


(a) *Chemical name*. Polyoxypropylene-polyoxyethylene glycol nonionic block polymer.

(b) *Specifications*. (1) Molecular weight range: 2,850 to 3,150.

(2) Hydroxyl number: 35.7 to 39.4.

(3) Cloud point (10 percent solution): 42° C.–46° C.

(4) Structural formula:



(c) *Sponsor*. (1) See No. 000069 in § 510.600(c) of this chapter for the sponsor of the usage provided by paragraph (d)(1) of this section.

(2) See No. 000069 in § 510.600(c) of this chapter for sponsor of usage provided by paragraph (d)(3) of this section.

(3) See No. 036904 in § 510.600(c) of this chapter for sponsor of usage provided by paragraph (d)(2) of this section.

(4) See No. 017800 in § 510.600(c) of this chapter for sponsor of the usage pro-

vided by paragraph (d)(4) of this section.

(d) *Conditions of use*. (1) For treatment of legume (alfalfa, clover) bloat in cattle. Administer as a drench at the rate of 25 grams for animals up to 500 pounds and 50 grams for animals over 500 pounds of body weight.

(2) For control of legume (alfalfa, clover) bloat in cattle. Administer, in molasses block containing 6.6 percent poloxalene, at the rate of 0.8 oz. of block (1.5 grams poloxalene) per 100 lbs. of body weight per day.

(3) For prevention of legume (alfalfa, clover) and wheat pasture bloat in cattle. A 53-percent poloxalene top dressing on individual rations of ground feed. Dosage is 1 gram of poloxalene per 100 pounds of body weight daily. If bloating conditions are severe, the dose is doubled. Treatment should be started 2 to 3 days before exposure to bloat-producing conditions. Repeat use of the drug if animals are exposed to bloat-producing conditions for more than 12 hours after the last treatment. Do not exceed the double dose in any 24-hour period.

(4) For control of legume (alfalfa, clover) and wheat pasture bloat in cattle. Administer in molasses block containing 6.6 percent poloxalene, at the rate of 0.8 ounce of block (1.5 grams of poloxalene) per 100 pounds of body weight per day. Provide access to blocks at least 7 days before exposure to bloat-producing conditions.

[40 FR 13838, Mar. 27, 1975, as amended at 40 FR 39857, Aug. 29, 1975; 42 FR 41854, Aug. 19, 1977; 50 FR 5385, Feb. 8, 1985; 54 FR 33501, Aug. 15, 1989; 56 FR 50653, Oct. 8, 1991; 58 FR 26523, May 4, 1993; 60 FR 55659, Nov. 2, 1995]

§ 520.1846 Polyoxyethylene (23) lauryl ether blocks.

(a) *Specifications*. Each molasses-based block contains 2.2 percent polyoxyethylene (23) lauryl ether.

(b) *Sponsor*. See No. 050112 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*. 2 grams of polyoxyethylene (23) lauryl ether per 100 kilograms of body weight per day (1 pound of block per 500 kilogram (1,100 pound) animal per day).

(2) *Indications for use*. For reduction of the incidence of bloat (alfalfa and clover) in pastured cattle.

§ 520.1870

(3) *Limitations.* Administer free-choice to beef cattle and nonlactating dairy cattle only. Initially, provide one block per five head of cattle. Start treatment 10 to 14 days before exposure to bloat-producing pastures. Do not allow cattle access to other sources of salt while being fed this product. Do not feed this product to animals without adequate forage/roughage consumption.

[50 FR 48189, Nov. 22, 1985, as amended at 56 FR 9841, Mar. 8, 1991]

§ 520.1870 Praziquantel tablets.

(a) *Specifications.* Each dog tablet contains 34 milligrams (mg) of praziquantel; each cat tablet contains 11.5 or 23 mg of praziquantel.

(b) *Sponsor.* See No. 000859 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Dogs—(i) Indications for use.* For removal of canine cestodes *Dipylidium caninum* and *Taenia pisiformis*. If labeled for use by or on the order of a licensed veterinarian, for removal of the canine cestode *Echinococcus granulosus*, and for removal and control of the canine cestode *Echinococcus multilocularis*.

(ii) *Dosage.* Dogs 5 pounds and under, ½ tablet (17 mg); 6 to 10 pounds, 1 tablet (34 mg); 11 to 15 pounds, 1½ tablets (51 mg); 16 to 30 pounds, 2 tablets (68 mg); 31 to 45 pounds, 3 tablets (102 mg); 46 to 60 pounds, 4 tablets (136 mg); over 60 pounds, 5 tablets maximum (170 mg).

(iii) *Limitations.* Administer directly by mouth or crumbled and in feed. Not intended for use in puppies less than 4 weeks of age. For over-the-counter (OTC) use: Consult your veterinarian before administering tablets to weak or debilitated animals, and for assistance in the diagnosis, treatment, and control of parasitism. For prescription use: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cats—(i) Indications for use.* For removal of feline cestodes *Dipylidium caninum* and *Taenia taeniaeformis*.

(ii) *Dosage.* Cats 4 pounds and under, 11.5 mg; 5 to 11 pounds, 23 mg; over 11 pounds, 34.5 mg.

(iii) *Limitations.* Administer directly by mouth or crumbled and in feed. Not intended for use in kittens less than 6 weeks of age. For OTC use: Consult

21 CFR Ch. I (4–1–01 Edition)

your veterinarian before administering tablets to weak or debilitated animals, and for assistance in the diagnosis, treatment, and control of parasitism.

[46 FR 60570, Dec. 11, 1981, as amended at 47 FR 26377, June 18, 1982; 55 FR 2234, Jan. 23, 1990; 58 FR 7864, Feb. 10, 1993; 58 FR 42853, Aug. 12, 1993]

§ 520.1871 Praziquantel/pyrantel pamoate tablets.

(a) *Specifications.* Each cat tablet contains 18.2 milligrams (mg) praziquantel with 72.6 mg pyrantel (as pyrantel pamoate).

(b) *Sponsor.* See 000859 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Cats—(i) Dosage.* 1.5 to 1.9 pounds, ¼ tablet; 2 to 3 pounds, ½ tablet; 4 to 8 pounds, 1 tablet; 9 to 12 pounds, 1 ½ tablets; 13 to 16 pounds, 2 tablets.

(ii) *Indications for use.* For removal of tapeworms (*Dipylidium caninum* and *Taenia taeniaeformis*), hookworms (*Ancylostoma tubaeforme*), and large roundworms (*Toxocara cati*) in cats and kittens.

(iii) *Limitations.* Not for use in kittens less than 1 month of age or weighing less than 1.5 pounds. May be given directly by mouth or in a small amount of food. Do not withhold food prior to or after treatment. If reinfection occurs, treatment may be repeated. Consult your veterinarian before giving to sick or pregnant animals. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(2) [Reserved]

[58 FR 58652, Nov. 3, 1993]

§ 520.1872 Praziquantel, pyrantel pamoate, and febantel tablets.

(a) *Specifications.* Each tablet contains either:

(1) Tablet No. 1: 22.7 milligrams praziquantel, 22.7 milligrams pyrantel base, and 113.4 milligrams febantel; or

(2) Tablet No. 2: 68 milligrams praziquantel, 68 milligrams pyrantel base, and 340.2 milligrams febantel.

(b) *Sponsor.* See 000859 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Dogs—(i) Amount.* Administer as a single dose directly by mouth or in a small amount of food as follows: