

§ 520.2043

21 CFR Ch. I (4-1-01 Edition)

2 to 4 weeks after first treatment regimen to determine the need for re-treatment. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[43 FR 52700, Nov. 14, 1978, as amended at 49 FR 22073, May 25, 1984; 57 FR 48163, Oct. 22, 1992; 58 FR 44611, Aug. 24, 1993]

§ 520.2043 Pyrantel pamoate suspension.

(a)(1) *Specifications.* Pyrantel pamoate suspension contains pyrantel pamoate equivalent to 50 milligrams of pyrantel base per milliliter.

(2) *Sponsors.* See Nos. 000069 and 059130 in § 510.600(c) of this chapter.

(3) *Conditions of use.* It is used in horses and ponies as follows:

(i) *Amount.* Equivalent of 3 milligrams pyrantel base per pound of body weight.

(ii) *Indications for use.* For the removal and control of infections from the following mature parasites: Large strongyles (*Strongylus vulgaris*, *Strongylus edentatus*, *Strongylus equinus*), small strongyles, pinworms (*Oxyuris*), and large roundworms (*Parascaris*).

(iii) *Limitations.* Administered as a single dose mixed with the usual grain ration, or by stomach tube, or by dose syringe. Not for use in horses and ponies to be slaughtered for food purposes. When the drug is for administration by stomach tube, it shall be labeled: "Federal law restricts this drug to use by or on the order of a licensed veterinarian." When the drug is not for administration by stomach tube, it shall be labeled: "Consult your veterinarian for assistance in the diagnosis, control, and treatment of parasitism."

(b)(1) *Specifications.* Pyrantel pamoate suspension contains pyrantel pamoate equivalent to 2.27 or 4.54 milligrams of pyrantel base per milliliter.

(2) *Sponsors.* See Nos. 000069, 011615, and 059130 for use of 2.27 and 4.54 milligrams per milliliter product. See No. 023851 for use of 4.54 milligrams per milliliter product.

(3) *Conditions of use.* It is used in puppies and dogs as follows:

(i) *Amount.* Equivalent of 2.27 milligrams of pyrantel base per pound of body weight.

(ii) *Indications for use.* For the removal of large roundworms (*Toxocara canis* and *Toxascarias leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*).

(iii) *Limitations.* Administer in the animal's feed bowl as a single dose by itself or mixed in a small quantity of food. Additional treatment may be required and should be confirmed by fecal examination within 2 to 4 weeks. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(4) *Conditions of use.* It is used in puppies and adult dogs and in lactating bitches after whelping as follows:

(i) *Amount.* Equivalent to 2.27 milligrams of pyrantel base per pound of body weight.

(ii) *Indications for use.* To prevent reinfections of *Toxocara canis*.

(iii) *Limitations.* Administer to puppies at 2, 3, 4, 6, 8, and 10 weeks of age. Administer to lactating bitches 2 to 3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

[42 FR 28534, June 3, 1977, as amended at 44 FR 36381, June 22 1979; 45 FR 10760, Feb. 19, 1980; 47 FR 16776, Apr. 20, 1982; 51 FR 11440, Apr. 3, 1986; 61 FR 29477, June 11, 1996; 61 FR 63712, Dec. 2, 1996; 62 FR 37712, July 15, 1997; 63 FR 39727, July 24, 1998; 63 FR 45945, Aug. 28, 1998]

§ 520.2044 Pyrantel pamoate paste.

(a) *Specifications.* Each milliliter of paste contains 180 milligrams of pyrantel base (as pyrantel pamoate).

(b) *Sponsor.* See 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use.* It is used in horses and ponies as follows:

(1) *Amount.* Equivalent of 3 milligrams pyrantel base per pound of body weight.

(2) *Indications for use.* For removal and control of infections from the following mature parasites: large strongyles (*Strongylus vulgaris*, *S. edentatus*, *S. equinus*); small strongyles; pinworms (*Oxyuris equi*); and large roundworms (*Parascaris equorum*).

(3) *Limitations.* Administer as single dose by depositing paste on dorsum of