

Food and Drug Administration, HHS

§ 520.2481

weight in a single dose. Dosage may be repeated in 21 days.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

§ 520.2460b Ticarbodine capsules.

(a) *Specifications.* Each capsule contains 90, 225, 450, or 900 milligrams of ticarbodine.

(b) *Sponsor.* See No. 000986 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in dogs for removal of roundworms (*Toxocara canis*), hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*), and tapeworms (*Dipylidium caninum* and *Taenia pisiformis*).

(2) Dosage is administered orally as a single dose at 45 milligrams per lb. of body weight. Dosage may be repeated at 21-day intervals.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

§ 520.2473 Tioxidazole oral dosage forms.

§ 520.2473a Tioxidazole granules.

(a) *Specifications.* Each gram of granules contains 200 milligrams of tioxidazole.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Horses—(i) Amount.* 5 milligrams per pound of body weight as a single dose.

(ii) *Indications for use.* Removal of mature large strongyles (*Strongylus edentatus*, *S. equinus*, and *S. vulgaris*), mature ascarids (*Parascaris equorum*), mature and immature (4th larval stage) pinworms (*Oxyuris equi*), and mature small strongyles (*Triodontophorus* spp.)

(iii) *Limitations.* For administration with feed: Sprinkle required amount of granules on a small amount of the usual grain ration and mix. Prepare for each horse individually. Withholding of feed or water not necessary. Not for use in horses intended for food. The reproductive safety of tioxidazole in breeding animals has not been determined. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. It is rec-

ommended that this drug be administered with caution to sick or debilitated horses.

(2) [Reserved]

[50 FR 52772, Dec. 26, 1985; 51 FR 2693, Jan. 21, 1986, as amended at 52 FR 7832, Mar. 13, 1987]

§ 520.2473b Tioxidazole paste.

(a) *Specifications.* Each plastic syringe contains 6.25 grams of tioxidazole.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Horses—(i) Amount.* 5 milligrams of tioxidazole per pound of body weight as a single dose.

(ii) *Indications for use.* Removal of mature large strongyles (*Strongylus edentatus*, *S. equinus*, and *S. vulgaris*), mature ascarids (*Parascaris equorum*), mature and immature (4th larval stage) pinworms (*Oxyuris equi*), and mature small strongyles (*Triodontophorus* spp.).

(iii) *Limitations.* Administer orally by inserting the nozzle of the syringe through the space between front and back teeth and deposit the required dose on the base of the tongue. Before dosing, make sure the horse's mouth contains no feed. Not for use in horses intended for food. The reproductive safety of tioxidazole in breeding animals has not been determined. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. It is recommended that this drug be administered with caution to sick or debilitated horses.

(2) [Reserved]

[52 FR 43059, Nov. 9, 1987]

§ 520.2481 Triamcinolone acetonide tablets.

(a) *Specifications.* Each tablet contains either 0.5 milligram or 1.5 milligrams of the drug.

(b) *Sponsor.* See Nos. 000010 and 053501 in § 510.600(c) of this chapter.

(c) *NAS/NRC status.* The conditions of use specified in this section are NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.