

§ 522.2220

21 CFR Ch. I (4-1-01 Edition)

(2) *Indications for use.* Treatment of diarrhea caused or complicated by *E. coli* (colibacillosis).

(3) *Limitations.* Administer as the sodium salt of sulfachlorpyridazine intravenously in aqueous solution for 1 to 5 days in divided doses twice daily; treated calves must not be slaughtered for food during treatment or for 5 days after the last treatment.

[40 FR 13858, Mar. 27, 1975, as amended at 50 FR 41489, Oct. 11, 1985]

§ 522.2220 Sulfadimethoxine injection.

(a)(1) *Specifications.* Sulfadimethoxine injection containing 400 milligrams per milliliter.

(2) *Sponsor.* (i) See No. 000069 in § 510.600(c) of this chapter for conditions of use as in paragraphs (a)(3)(i) through (a)(3)(iii) of this section.

(ii) See No. 057561 for conditions of use as in paragraph (a)(3) of this section.

(iii) See No. 059130 for use as in paragraph (a)(3)(iii) of this section.

(3) *Conditions of use.* (i) It is used or intended for use in dogs and cats as follows:

(a) For the treatment of respiratory, genitourinary tract, enteric, and soft tissue infections when caused by Streptococci, Staphylococci, Escherichia, Salmonella, Klebsiella, Proteus, or Shigella organisms sensitive to sulfadimethoxine, and in the treatment of canine bacterial enteritis associated with coccidiosis and canine Salmonellosis.

(b) It is administered by intravenous or subcutaneous injection at an initial dose of 55 milligrams per kilogram of body weight followed by 27.5 milligrams per kilogram of body weight every 24 hours.

(c) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(ii) It is used or intended for use in horses as follows:

(a) For the treatment of respiratory disease caused by Streptococcus equi (strangles).

(b) It is administered by intravenous injection at an initial dose of 55 milligrams per kilogram of body weight followed by 27.5 milligrams per kilogram of body weight every 24 hours until the patient is asymptomatic for 48 hours.

(c) Not for use in horses intended for food.

(d) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(iii) It is used or intended for use in cattle as follows:

(a) For the treatment of shipping fever complex, bacterial pneumonia, calf diphtheria, and foot-rot.

(b) It is administered by intravenous injection at an initial dose of 25 milligrams per pound of body weight followed by 12.5 milligrams per pound of body weight every 24 hours until the animal is asymptomatic for 48 hours.

(c) Milk taken from animals during treatment and for 60 hours (5 milkings) after the latest treatment must not be used for food. Do not administer within 5 days of slaughter.

(d) Tissue damage may result from perivascular infiltration.

(b) [Reserved]

(c)(1) *Specifications.* Sulfadimethoxine containing 100 milligrams per milliliter.

(2) *Sponsor.* See No. 000010 in § 510.600(c) of this chapter.

(3) *Conditions of use.* (i) It is used or intended for use in the treatment of sulfadimethoxine-susceptible bacterial infections in dogs.

(ii) It is administered by subcutaneous, intramuscular, or intravenous injection at an initial dose of 25 milligrams per pound of body weight followed by 12.5 milligrams per pound of body weight every 24 hours thereafter. Continue treatment until the animal is free from symptoms for 48 hours.

(iii) For use by or on the order of a licensed veterinarian.

(d) *Related tolerances.* See § 556.640 of this chapter.

[40 FR 13858, Mar. 27, 1975, as amended at 40 FR 34112, Aug. 14, 1975; 40 FR 42007, Sept. 10, 1975; 50 FR 254, Jan. 3, 1985; 53 FR 40728, Oct. 18, 1988; 54 FR 30205, July 19, 1989; 58 FR 38972, July 21, 1993; 59 FR 56000, Nov. 10, 1994; 61 FR 4875, Feb. 9, 1996; 62 FR 23128, Apr. 29, 1997; 62 FR 35076, June 30, 1997]

§ 522.2240 Sulfaethoxy-pyridazine.

(a) *Chemical name.* N¹-(6-Ethoxy-3-pyridazinyl) sulfanilamide.

(b) *Specifications.* Melting point range of 180° C to 186° C.

Food and Drug Administration, HHS

§ 522.2340

(c) *Sponsor.* See No. 010042 in § 510.600(c) of this chapter.

(d) *Related tolerances.* See § 556.650 of this chapter.

(e) *Conditions of use.* It is used for injection into cattle as follows:

(1) *Amount.* 2.5 grams per 100 pounds of body weight per day.

(2) *Indications for use.* Treatment of respiratory infection (pneumonia, shipping fever), foot rot, calf scours; as adjunctive therapy in septicemia accompanying mastitis and metritis.

(3) *Limitations.* Administer intravenously for not more than 4 days; or first treatment may be followed by 3 days of treatment with sulfaethoxyypyridazine in drinking water, feed, or tablet in accordance with § 558.579(e) or §§ 520.2240a(e) and 520.2240b(e) of this chapter; as sodium sulfaethoxyypyridazine; do not treat within 16 days of slaughter; as sole source of sulfonamide; milk that has been taken from animals during treatment and for 72 hours (6 milkings) after the latest treatment must not be used for food; for use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 41 FR 11011, Mar. 15, 1976]

§ 522.2260 Sulfamethazine injectable solution.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 250 milligrams of sulfamethazine sodium.

(b) *Sponsor.* See No. 010042 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.670 of this chapter.

(d) *NAS/NRC status.* The conditions of use are NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(e) *Conditions of use—(1) Amount.* 20 milliliters for each 50 pounds of body weight (100 milligrams per pound) initially, 20 milliliters per 100 pounds of body weight (50 milligrams per pound) daily thereafter for cattle.

(2) *Indications for use.* For cattle for treatment of bacterial pneumonia and bovine respiratory disease complex (shipping fever complex) (*Pasteurella* spp.), colibacillosis (bacterial scours)

(*Escherichia coli*), necrotic pododermatitis (foot rot) (*Fusobacterium necrophorum*), calf diphtheria (*Fusobacterium necrophorum*), acute mastitis and acute metritis (*Streptococcus* spp.) when caused by one or more pathogenic organisms sensitive to sulfamethazine.

(3) *Limitations.* For intravenous use only. Not for use in lactating dairy animals. Withdraw medication from cattle 10 days prior to slaughter for food. If symptoms persist for 2 or 3 days, consult a veterinarian. Adequate water intake is important for animals treated with sulfonamides. Treatment should continue 24 to 48 hours beyond the remission of disease symptoms, but not to exceed a total of 5 consecutive days.

[46 FR 62055, Dec. 22, 1981]

§ 522.2340 Sulfomyxin.

(a) *Specifications.* Sulfomyxin for injection is sterile. It is derived from the antibiotic substance produced by the growth of *Bacillus polymyxa* or is the same substance produced by any other means.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Special considerations.* The quantities of antibiotic in paragraph (e) of this section refer to the activity of the appropriate standard.

(d) *Related tolerances.* See § 556.700 of this chapter.

(e) *Conditions of use.* (1) It is used or intended for use in chickens and turkeys as an aid in the treatment of disease caused or complicated by *E. coli*, such as colibacillosis and complicated chronic respiratory disease.

(2) It is administered by subcutaneous injection as follows:

Age of birds in days	Antibiotic activity	
	Chickens (units)	Turkeys (units)
1 to 14	12,500	12,500
15 to 28	25,000	25,000
29 to 63	50,000	50,000
Over 63	50,000	100,000

(3) A second injection may be given 3 days later if symptoms persist.

(4) Not for use in laying hens; do not treat chickens within 5 days of slaughter; do not treat turkeys within 7 days of slaughter.