

by or on the order of a licensed veterinarian.

[59 FR 5105, Feb. 3, 1994]

§ 524.1484i Neomycin sulfate, hydrocortisone acetate, sterile ointment.

(a) *Specifications.* The drug contains 5 milligrams of neomycin sulfate, equivalent to 3.5 milligrams of neomycin base, and 5 milligrams of hydrocortisone acetate in each gram of ointment.¹

(b) *Sponsor.* No. 000009 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) *Amount.* Apply three or four times daily into the conjunctival sac. With improvement, frequency may be reduced to two or three times daily. For treatment of ear canker and other inflammatory conditions of the external ear canal, fill external ear canal one to three times daily.¹

(2) *Indications for use.* For treating infections, allergic, and traumatic keratitis, conjunctivitis, acute otitis externa and, to a lesser degree, chronic otitis externa in dogs and cats.¹

(3) *Limitations.* All topical ophthalmic preparations containing corticosteroids, with or without an antimicrobial agent, are contraindicated in the initial treatment of corneal ulcers. They should not be used until infection is under control and corneal regeneration is well underway. Incomplete response or exacerbation of corticosteroid responsive lesions may be due to the presence of nonsusceptible organisms or to prolonged use on antibiotic-containing preparations resulting in overgrowth of nonsusceptible organisms, particularly *Monilia*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[43 FR 40456, Sept. 12, 1978]

§ 524.1484j [Reserved]

§ 524.1484k Neomycin sulfate, prednisolone, tetracaine, and squalane topical-otic suspension.

(a) *Specifications.* Each milliliter of suspension contains 5 milligrams neomycin sulfate (equivalent to 3.5 milligrams neomycin base), 2 milligrams prednisolone, 5 milligrams tetracaine, and 0.25 milliliter squalane.

(b) *Sponsor.* See 017030 in § 510.600(c) of this chapter.

(c) *Conditions of use—*(1) *Amount.* 2 to 3 applications daily or as needed.

(2) *Indications for use.* Indicated for use in dogs and cats for treating acute otitis externa and as adjunctive therapy in management of chronic otitis externa. The product may also be used for treating moist dermatitis in dogs.

(3) *Limitations.* Tetracaine and neomycin have the potential to sensitize. If signs of irritation or sensitivity develop, discontinue use. Prolonged use of this product may result in overgrowth of nonsusceptible organisms. If new infections due to bacteria or fungi appear during therapy, appropriate measures should be taken. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[48 FR 5265, Feb. 4, 1983; 48 FR 8055, Feb. 25, 1983]

§ 524.1580 Nitrofurazone ophthalmic and topical dosage forms.

§ 524.1580a [Reserved]

§ 524.1580b Nitrofurazone ointment.

(a) *Specifications.* The drug contains 0.2 percent nitrofurazone in a water-soluble base.

(b) *Sponsor.* For use on dogs, cats, or horses, see Nos. 000010, 000857, 000864, 000069, 050749, 023851, 051259, and 061133 in § 510.600(c) of this chapter. For use on dogs and horses, see No. 017135 in § 510.600(c) of this chapter. For use on horses, see No. 017153 in § 510.600(c) of this chapter.

(c) *Conditions of use—*(1) *Indications for use.* For prevention or treatment of surface bacterial infections of wounds, burns, and cutaneous ulcers of dogs, cats, or horses.¹

(2) *Limitations.* Apply directly on the lesion with a spatula or first place on a piece of gauze. Use of a bandage is optional. The preparation should remain on the lesion for at least 24 hours. The dressing may be changed several times daily or left on the lesion for a longer period. For use only on dogs, cats, and horses (not for food use). In case of deep or puncture wounds or serious burns, use only as recommended by veterinarian. If redness, irritation, or

§ 524.1580c

swelling persists or increases, discontinue use; consult veterinarian.¹

[46 FR 43402, June 27, 1980, as amended at 49 FR 6476, Feb. 22, 1984; 50 FR 49373, Dec. 2, 1985; 52 FR 18691, May 19, 1987; 53 FR 32610, Aug. 26, 1988; 53 FR 40728, Oct. 18, 1988; 54 FR 29544, July 13, 1989; 54 FR 37097, Sept. 7, 1989; 55 FR 8462, Mar. 8, 1990; 55 FR 20455, May 17, 1990; 56 FR 37473, Aug. 7, 1991; 56 FR 50653, Oct. 8, 1991; 59 FR 33197, June 28, 1994; 60 FR 55659, Nov. 2, 1995; 62 FR 35077, June 30, 1997; 66 FR 14074, Mar. 9, 2001]

§ 524.1580c Nitrofurazone soluble powder.

(a) *Specifications.* The drug contains 0.2 percent nitrofurazone in a water-soluble base.

(b) *Sponsor.* See Nos. 000010, 000069, and 050749 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Amount.* Apply several times daily to the lesion or affected area from the plastic squeeze bottle.

(2) *Indications for use.* For prevention or treatment of surface bacterial infections of wounds, burns, skin ulcers, and abscesses after incision.¹

(3) *Limitations.* In case of deep or puncture wounds or serious burns, use only as recommended by veterinarian. If redness, irritation, or swelling persists or increases, discontinue use; consult veterinarian. For use only on dogs, cats, and horses (not for food use).¹

[45 FR 43402, June 27, 1980, as amended at 47 FR 43368, Oct. 1, 1982; 48 FR 28984, June 24, 1983; 53 FR 40728, Oct. 18, 1988; 54 FR 30542, July 21, 1989; 56 FR 50653, Oct. 8, 1991; 59 FR 33197, June 28, 1994; 60 FR 55659, Nov. 2, 1995; 62 FR 35077, June 30, 1997]

§ 524.1580d [Reserved]

§ 524.1580e Nitrofurazone ointment with butacaine sulfate.

(a) *Specifications.* The drug contains 0.2 percent nitrofurazone and 0.5 percent butacaine sulfate in a water-soluble base.

(b) *Sponsor.* See No. 051259 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Indications for use.* For prevention or treatment of surface bacterial infections of ears, wounds, burns, and cutaneous ulcers of dogs, cats, and horses.¹

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these

21 CFR Ch. I (4–1–01 Edition)

(2) *Limitations.* Apply directly on the lesion with a spatula or first place on a piece of gauze. Use of a bandage is optional. The preparation should remain on the lesion for at least 24 hours. The dressing may be changed several times daily or left on the lesion for a longer period. For use only on dogs, cats, and horses (not for food use). In case of deep or puncture wounds or serious burns, use only as recommended by a veterinarian. If redness, irritation, or swelling persists or increases, discontinue use; consult veterinarian.¹

[49 FR 9417, Mar. 13, 1984]

§ 524.1600 Nystatin ophthalmic and topical dosage forms.

§ 524.1600a Nystatin, neomycin, thiostrepton, and triamcinolone acetone ointment.

(a) *Specifications.* Each milliliter of petrolatum base or each gram of vanishing cream base ointment contains: 100,000 units of nystatin; neomycin sulfate equivalent to 2.5 milligrams of neomycin base; 2,500 units of thiostrepton; and 1.0 milligram of triamcinolone acetone.

(b) *Sponsors.* For petrolatum base ointments see 000031, 000069, 000332, 025463, 051259, and 053501 in § 510.600(c) of this chapter. For vanishing cream base ointments see 051259 and 053501.

(c) *Conditions of use—(1) Amount.* (i) For topical dermatological use: Clean affected areas and remove any encrusted discharge or exudate, and apply sparingly either ointment in a thin film.

(ii) For otic use: Clean ear canal of impacted cerumen, remove any foreign bodies such as grass awns and ticks, and instill three to five drops of petrolatum base ointment. Preliminary use of a local anesthetic may be advisable.

(iii) For infected anal glands and cystic areas: Drain gland or cyst and fill with petrolatum base ointment.

(2) *Indications for use.* (i) Topically: Use either ointment in dogs and cats for anti-inflammatory, antipruritic,

uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.