

(b) *Sponsor*. See No. 017030 in § 510.600(c) of this chapter.

(c) *Conditions of use. Dogs and cats.* (1) *Amount*. Apply 4 to 6 times daily to the affected eye for the first 72 hours depending upon the severity of the condition. Continue treatment for 48 hours after the eye appears normal.

(2) *Indications for use*. Treatment of bacterial conjunctivitis and ocular inflammation caused by organisms susceptible to chloramphenicol.

(3) *Limitations*. Therapy for cats should not exceed 7 days, prolonged use in cats may produce blood dyscrasia. As with other antibiotics, prolonged use may result in overgrowth of non-susceptible organisms. If superinfection occurs or if clinical improvement is not noted within a reasonable period, discontinue use and institute appropriate therapy. All topical ophthalmic preparations containing corticosteroids, with or without an antimicrobial agent, are contraindicated in the initial treatment of corneal ulcers. They should not be used until the infection is under control and corneal regeneration is well underway. Chloramphenicol products must not be used in meat-, egg-, or milk-producing animals. The length of time that residues persist in milk or tissues has not been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37334, Aug. 18, 1992]

#### § 524.402 Chlorhexidine diacetate ointment.

(a) *Specifications*. The product contains 1 percent of chlorhexidine diacetate in an ointment base.

(b) *Sponsor*. See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used as a topical antiseptic ointment for surface wounds on dogs, cats, and horses.<sup>1</sup>

(2) The wound area is carefully cleansed and the drug is applied daily.<sup>1</sup>

<sup>1</sup>These conditions are NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(3) The drug is not to be used in horses intended for use as food.<sup>1</sup>

#### § 524.450 Clotrimazole cream.

(a) *Specifications*. Each gram of cream contains 10 milligrams of clotrimazole.

(b) *Sponsor*. See 000859 in § 510.600(c).

(c) *Conditions of use*—(1) *Amount*. Apply ¼-inch ribbon of cream per square inch of lesion once daily for 2 to 4 weeks.

(2) *Indications for use*. For the treatment of fungal infections of dogs and cats caused by *Microsporium canis* and *Trichophyton mentagrophytes*.

(3) *Limitations*. Wash hands thoroughly after use to avoid spread of infection. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 48128, July 18, 1980]

#### § 524.463 Copper naphthenate solution.

(a) *Specifications*. The drug contains 37.5 percent copper naphthenate in a suitable solvent.

(b) *Sponsor*. See Nos. 000856 and 017135 in § 510.600(c) of this chapter.

(c) *Conditions of use*—*Horses and ponies*—(1) *Amount*. Apply daily to affected hooves until fully healed.

(2) *Indications for use*. As an aid in treating horses and ponies for thrush caused by organisms susceptible to copper naphthenate.

(3) *Limitations*. Use on horses and ponies only. Remove debris and necrotic material before applying. Avoid contact around eyes. Do not use on animals that are raised for food production. Do not contaminate feed. Do not allow runoff of excess drug into hair because contact with the drug may cause some hair loss.

[47 FR 4250, Jan. 29, 1982]

#### § 524.520 Cuprimyxin cream.

(a) *Specifications*. The drug contains 0.5 percent cuprimyxin (6-methoxy-1-phenazinol 5, 10-dioxide, cupric complex) in an aqueous cream base.

(b) *Sponsor*. See No. 000004 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) Cuprimyxin is a broad spectrum antibacterial and antifungal cream for the topical treatment of superficial infections in