

§ 524.770 Doramectin.

(a) *Specifications.* Each milliliter of solution contains 5 milligrams of doramectin.

(b) *Sponsor.* See 000069 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.225 of this chapter.

(d) *Conditions of use—Cattle—(1) Amount.* 5 milligrams per 10 kilograms (5 milligrams per 22 pounds).

(2) *Indications for use.* For treatment and control of gastrointestinal roundworms, lungworms, eyeworms, grubs, biting and sucking lice, horn flies, and mange mites. To control infections and to protect from reinfection with *Cooperia oncophora* and *Dictyocaulus viviparus* for 21 days, *Ostertagia ostertagi*, *C. punctata*, and *Oesophagostomum radiatum* for 28 days, and *Haemonchus placei* for 35 days after treatment.

(3) *Limitations.* Administer as a single dose. Do not slaughter cattle within 45 days of latest treatment. Not for use in female dairy cattle 20 months of age or older. Do not use in calves to be processed for veal. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[62 FR 65753, Dec. 16, 1997, as amended at 63 FR 68183, Dec. 10, 1998; 64 FR 49082, Sept. 10, 1999]

§ 524.802 Enrofloxacin, silver sulfadiazine emulsion.

(a) *Specifications.* Each milliliter contains 5 milligrams (mg) enrofloxacin and 10 mg silver sulfadiazine.

(b) *Sponsor.* See No. 000859 in § 510.600(c) of this chapter.

(c) *Conditions of use—Dogs—(1) Amount.* 5 to 10 drops for dogs weighing 35 pounds (lb) or less and 10 to 15 drops for dogs weighing more than 35 lb; applied twice daily for up to 14 days.

(2) *Indications for use.* For the treatment of otitis externa in dogs.

(3) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian. Federal law prohibits the extra-label use of this drug in food-producing animals.

[65 FR 66620, Nov. 7, 2000]

§ 524.814 Eprinomectin.

(a) *Specifications.* Each milliliter contains 5 milligrams of eprinomectin.

(b) *Sponsor.* See No. 000006 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.227 of this chapter.

(d) *Conditions of use—(1) Amount.* One milliliter (5 milligrams) per 10 kilograms of body weight (500 micrograms per kilogram).

(2) *Indications for use.* The drug is used in beef and dairy cattle for treatment and control of gastrointestinal roundworms (*Haemonchus placei* (adult and L4), *Ostertagia ostertagi* (adult and L4, including inhibited L4), *Trichostrongylus axei* (adult and L4), *T. colubriformis* (adult and L4), *T. longispicularis* (adult), *Cooperia oncophora* (adult and L4), *C. punctata* (adult and L4), *C. surnabada* (adult and L4), *Nematodirus helvetianus* (adult and L4), *Bunostomum phlebotomum* (adult and L4), *Oesophagostomum radiatum* (adult and L4), *Strongyloides papillosus* (adults), *Trichuris* spp. (adults)); lungworms (*Dictyocaulus viviparus*, adult and L4); cattle grubs (all parasitic stages *Hypoderma lineatum*, *H. bovis*); lice (*Damalinia bovis*, *Linognathus vituli*, *Haematopinus eurysternus*, *Solenopotes capillatus*); mange mites (*Chorioptes bovis*, *Sarcoptes scabiei*); and horn flies (*Haematobia irritans*). Controls and protects from reinfection of *D. viviparus* for 21 days after treatment and *H. irritans* for 7 days after treatment.

(3) *Limitations.* Apply topically along backbone from withers to tailhead. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[62 FR 33997, June 24, 1997, as amended at 63 FR 59715, Nov. 5, 1998]

§ 524.900 Famphur.

(a) *Chemical name.* *O,O*- Dimethyl *O*-[*p*-(dimethylsulfamoyl)phenyl] phosphorothioate.

(b) *Specifications.* The drug is in liquid form containing 13.2 percent famphur.

(c) *Sponsor.* See Nos. 000061 and 060594 in § 510.600(c) of this chapter.

(d) *Special considerations.* Do not use on animals simultaneously or within a few days before or after treatment with

§ 524.920

21 CFR Ch. I (4-1-01 Edition)

or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(e) *Related tolerances.* See §556.273 of this chapter.

(f) *Conditions of use.* (1) The drug is used as a pour-on formulation for the control of cattle grubs and to reduce cattle lice infestations.

(2) It is used at the rate of 1 ounce per 200 pounds body weight, not to exceed a total dosage of 4 ounces, applied from the shoulder to the tail head as a single treatment. It is applied as soon as possible after heel fly activity ceases. Do not use on lactating dairy cows or dry dairy cows within 21 days of freshening, calves less than 3 months old, animals stressed from castration, over-excitement or dehorning, sick or convalescent animals. Animals may become dehydrated and under stress following shipment. Do not treat until they are in good condition. Brahman and Brahman crossbreeds are less tolerant of cholinesterase-inhibiting insecticides than other breeds. Do not treat Brahman bulls.

(3) Do not slaughter within 35 days after treatment. Swine should be eliminated from area where run-off occurs.

[40 FR 13873, Mar. 27, 1975, as amended at 49 FR 34352, Aug. 30, 1984; 57 FR 7652, Mar. 4, 1992; 59 FR 28769, June 3, 1994; 62 FR 55161, Oct. 23, 1997; 62 FR 61626, Nov. 19, 1997]

§ 524.920 Fenthion.

(a) *Chemical name.* O,O-Dimethyl O-[4-(methylthio)-*m*-tolyl] phosphorothioate.

(b) *Specifications.* (1) The drug is in a liquid form containing 3 percent of fenthion.

(2) *Sponsor.* See No. 000859 in §510.600(c) of this chapter.

(3) *Special considerations.* Do not use on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(4) *Related tolerances.* See 40 CFR 180.214.

(5) *Conditions of use.* (i) The drug is used as a pour-on formulation for the control of grubs and lice in beef and nonlactating cattle.

(ii) It is used at the rate of one-half fluid ounce per 100 pounds of body weight placed on the backline of the animal. Only one application per sea-

son should be made for grub control and this will also provide initial control of lice. A second application for lice control may be made if animals become reinfested, but no sooner than 35 days after the first treatment. Proper timing of treatment is important for grub control; cattle should be treated as soon as possible after heel-fly activity ceases. Cattle should not be slaughtered within 35 days following a single treatment. If a second application is made for lice control, cattle should not be slaughtered within 45 days of the second treatment. The drug must not be used within 28 days of freshening of dairy cattle. If freshening should occur within 28 days after treatment, do not use milk as human food for the balance of the 28-day interval. Do not treat lactating dairy cattle; calves less than 3 months old; or sick, convalescent, or stressed livestock. Do not treat cattle for 10 days before or after shipping, weaning, or dehorning or after exposure to contagious infectious diseases.

(c) *Specifications.* (1) The drug is in a liquid form containing 20 percent fenthion.

(2) *Sponsor.* See No. 000859 in §510.600(c) of this chapter.

(3) *Special considerations.* Do not use on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(4) *Related tolerances.* See 40 CFR 180.214.

(5) *Conditions of use.* (i) The drug is used for control of cattle grubs and as an aid in controlling lice on beef cattle and on dairy cattle not of breeding age.

(ii) It is applied as a single application placed on the backline of animals as follows:

Weight of animal	Dosage (milliliters)
150 to 300 lb	4
301 to 600 lb	8
601 to 900 lb	12
901 to 1,200 lb	16
Over 1,200 lb	20

For most effective results, cattle should be treated as soon as possible after heel-fly activity ceases. Host-parasite reactions such as bloat, salivation, staggering and paralysis may