

(b) *Sponsors.* See Nos. 000074, 000209, 010019, 012164, and 059258 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*—(i) *Horses:* For induction of surgical anesthesia: 3 to 5 percent isoflurane (with oxygen) for 5 to 10 minutes. For maintenance of surgical anesthesia: 1.5 to 1.8 percent isoflurane (with oxygen).

(ii) *Dogs:* For induction of surgical anesthesia: 2 to 2.5 percent isoflurane (with oxygen) for 5 to 10 minutes. For maintenance of surgical anesthesia: 1.5 to 1.8 percent isoflurane (with oxygen).

(2) *Indications for use.* For induction and maintenance of general anesthesia in horses and dogs.

(3) *Limitations.* Administer by inhalation; not for use in horses or dogs sensitive to halogenated agents; increasing depth of anesthesia may increase hypotension and respiratory depression; use less than usual amounts of nondepolarizing relaxants; use with vaporizers producing predictable percentage concentrations; not for use in horses intended for food; Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[51 FR 594, Jan. 7, 1986, as amended at 54 FR 23472, June 1, 1989; 58 FR 17346, Apr. 2, 1993; 59 FR 44315, Aug. 29, 1994; 60 FR 40456, Aug. 9, 1995; 63 FR 8122, Feb. 18, 1998; 63 FR 24106, May 1, 1998]

#### § 529.1526 Nifurpirinol capsules.

(a) *Specifications.* Each capsule contains 3.8 or 7.6 milligrams of nifurpirinol.

(b) *Sponsor.* See No. 000074 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in treating aquarium fish for the control of columnaris disease caused by *Chondrococcus columnaris* susceptible to nifurpirinol.

(2) Use one 3.8 milligram nifurpirinol capsule for each 10 gallons of aquarium water. Empty the contents of the capsule directly into the water and stir briefly. Treat for at least 1 hour. If activated charcoal or carbon filtration is being used, disconnect during treatment, but maintain adequate aeration. Resume water filtration after 1 hour treatment. Usually a single treatment is sufficient. For aquariums with charcoal filters, nifurpirinol can be used once each 24 hours up to 3 consecutive

days, discontinuing filtration during treatment. If aquarium does not have charcoal filter, do not retreat within 5 days.

(3) Do not use in salt water aquariums.

(4) Do not use while egg bearers or live bearers are reproducing.

[40 FR 60052, Dec. 31, 1975, as amended at 47 FR 20758, May 14, 1982; 56 FR 43699, Sept. 4, 1991]

#### § 529.2150 Sevoflurane.

(a) *Specifications.* The drug is a clear, colorless, stable liquid containing no additives or chemical stabilizers.

(b) *Sponsor.* See No. 000074 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* For induction of surgical anesthesia: 5 to 7 percent sevoflurane with oxygen. For maintenance of surgical anesthesia: 3.7 to 4 percent sevoflurane with oxygen in the absence of premedication and 3.3 to 3.6 percent in the presence of premedication.

(2) *Indications for use.* For induction and maintenance of general anesthesia in dogs.

(3) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[64 FR 71640, Dec. 22, 1999]

#### § 529.2090 Salicylic acid.

(a) *Specifications.* (1) Each dose contains 0.55 grain of salicylic acid in a gum arabic and dextrin vehicle.

(2) Each dose is incorporated upon a device (teat dilator) suitable for insertion into and subsequent removal from the teat canal.

(b) *Sponsor.* See No. 045087 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used for the removal of scar tissue in the teat canal of milk-producing cows.

(2) The labeling bears directions to the user to:

(i) Treat lactating cows initially by inserting dosage and removal of the device;

(ii) Insert second dose and permit device to remain in canal until the next milking; and

(iii) Insert one dose following each milking for not more than 2 days.