

§ 558.145 Chlortetracycline, procaine penicillin, and sulfamethazine.

(a) *Approvals.* Type A medicated articles: (1) 20 grams of chlortetracycline per pound, 4.4 percent (20 grams) of sulfamethazine, and procaine penicillin equivalent in activity to 10 grams of penicillin per pound to 046573 and 063238 in § 510.600(c) of this chapter.

(2) 40 grams of chlortetracycline per pound, 8.8 percent of sulfamethazine, and penicillin procaine equivalent in activity to 20 grams of penicillin per pound to 063238 in § 510.600(c) of this chapter.

(b) *Specifications.* (1) The antibiotic substance refers to the antibiotic or feed-grade antibiotic.

(2) The antibiotic activities are expressed in terms of the appropriate antibiotic standards.

(3) Type C medicated feed contains in each ton, 100 grams of chlortetracycline, 50 grams of penicillin as procaine penicillin, and 100 grams of sulfamethazine.

(c) *Related tolerances.* See §§ 556.150, 556.510, and 556.670 of this chapter.

(d) *Conditions of use.* (1) It is administered to swine in a Type C feed for reduction of the incidence of cervical abscesses; treatment of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by *Salmonella choleraesuis* and vibronic dysentery); prevention of these diseases during times of stress; maintenance of weight gains in the presence of atrophic rhinitis; growth promotion and increased feed efficiency in swine weighing up to 75 pounds.

(2) Withdraw 15 days prior to slaughter.

[40 FR 13959, Mar. 27, 1975, as amended at 43 FR 19385, May 5, 1978; 47 FR 39814, Sept. 10, 1982; 48 FR 30615, July 5, 1983; 51 FR 7396, Mar. 3, 1986; 52 FR 2684, Jan. 26, 1987; 56 FR 14019, Apr. 5, 1991; 61 FR 18082, Apr. 24, 1996; 62 FR 14300, Mar. 26, 1997; 63 FR 27845, May 21, 1998]

§ 558.155 Chlortetracycline, sulfathiazole, penicillin.

(a) *Approvals.* Type A medicated articles: (1) 20 grams of chlortetracycline hydrochloride, 4.4 percent (20 grams) sulfathiazole, and procaine penicillin equivalent to 10 grams of penicillin per

pound to 000004 and 000010 in § 510.600(c) of this chapter.

(2) 40 grams of chlortetracycline hydrochloride, 8.8 percent (40 grams) sulfathiazole and procaine penicillin equivalent in activity to 20 grams of penicillin per pound to Nos. 000004 and 000010 in § 510.600(c) of this chapter.

(b) *Specifications.* (1) The antibiotic substance refers to the antibiotic or feed-grade antibiotic.

(2) The antibiotic activities are expressed in terms of the appropriate antibiotic standards.

(c) *Related tolerances.* See §§ 556.150, 556.510, and 556.690 of this chapter.

(d) *Conditions of use.* It is used for swine as follows:

(1) *Amount per ton.* Chlortetracycline, 100 grams plus penicillin, 50 grams plus sulfathiazole, 100 grams.

(2) *Indications for use.* For reduction of incidence of cervical abscesses. Treatment of bacterial enteritis (salmonellosis or necrotic enteritis caused by *Salmonella choleraesuis* and vibronic dysentery). Maintenance of weight gains in the presence of atrophic rhinitis. Swine 10 pounds of body weight to 6 weeks post-weaning: Increased rate of weight gain and improved feed efficiency. Swine 6 to 16 weeks post-weaning: Increased rate of weight gain.

(3) *Limitations.* For swine raised in confinement (dry-lot) or on limited pasture. Feed as sole ration. Withdraw 7 days prior to slaughter.

MINIMUM AMOUNT OF TYPE C FEED WHICH THE ANIMAL SHOULD CONSUME

Type of feed	Approximate body weight in pounds	Minimum desired daily feed intake in pounds
Prestarter (up to 6 weeks postweaning)	20	1
Starter (up to 6 weeks postweaning)	50	1½
Grower (6-16 weeks postweaning)	80	2
Finisher (6-16 weeks postweaning)	150	3

[40 FR 13959, Mar. 27, 1975, as amended at 51 FR 7397, Mar. 3, 1986; 51 FR 28547, Aug. 8, 1986; 52 FR 2684, Jan. 26, 1987; 61 FR 2415, Jan. 26, 1996; 62 FR 35077, June 30, 1997; 62 FR 67725, Dec. 30, 1997; 63 FR 11599, Mar. 10, 1998]