

**Food and Drug Administration, HHS**

**§ 558.254**

(B) *Limitations.* Feed continuously as sole ration. Not to be used in swine weighing more than 250 pounds.

(2) [Reserved]

[57 FR 38442, Aug. 25, 1992, as amended at 62 FR 63271, Nov. 28, 1997]

**§ 558.248 Erythromycin thiocyanate.**

(a) *Approvals.* Type A medicated articles: (1) 2.2 percent to 061133 in § 510.600(c) of this chapter for use as in paragraph (d) of this section.

(2) 5 and 10 percent to 061133 for use in paragraphs (d)(1)(i) and (ii) of this section.

(b) *Special considerations.* The levels of antibiotic are expressed in terms of erythromycin master standard. One gram of erythromycin thiocyanate is equivalent to 0.925 gram of erythromycin master standard.

(c) *Related tolerances.* See § 556.230 of this chapter.

(d) *Condition of use.* (1) It is used as follows:

Erythromycin thiocyanate in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) 4.6 to 18.5		Chickens; growth promotion and feed efficiency.		061133
(ii) 9.25 to 18.5		Turkeys; growth promotion and feed efficiency.	For turkeys not over 12 weeks of age.	061133
(iii) 9.25 to 64.75		Swine; increase in weight gain, improved feed efficiency in starter pigs (9.25 to 64.75) and grower-finishing pigs (9.25).	Starter ration for animals up to 35 lb body weight.	061133
(iv) 18.5		Laying chickens; aids in increasing egg production.		061133
(v) 92.5		1. Chickens; as an aid in the prevention of chronic respiratory disease during periods of stress. 2. Chickens; as an aid in the prevention of infectious coryza. 3. Turkeys; as an aid in the prevention of chronic respiratory disease during periods of stress.	Feed for 2 d before stress and 3 to 6 d after stress; withdraw 24 h before slaughter. Feed for 7 to 14 d; withdraw 24 h before slaughter. Feed for 2 d before stress and 3 to 6 d after stress.	061133
(vi) 185		1. Chickens; as an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease. 2. Turkeys; as an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease.	Feed for 5 to 8 d; do not use in birds producing eggs for food purposes; withdraw 48 h before slaughter. Feed for 5 to 8 d; do not use in birds producing eggs for food purposes.	061133

(2) In feed for feedlot beef cattle at 37 milligrams per head per day as an aid in stimulating growth and improving feed efficiency.

(3) Erythromycin thiocyanate may be used in accordance with the provisions of this section in the combinations provided as follows:

(i) Amprolium in accordance with § 558.55.

(ii) Amprolium and ethopabate in accordance with § 558.58.

(iii) Arsanilic acid in accordance with § 558.62.

(iv) Zoalene in accordance with § 558.680.

[41 FR 10999, Mar. 15, 1976, as amended at 45 FR 56799, Aug. 26, 1980; 49 FR 31281, Aug. 6, 1984; 51 FR 7397, Mar. 3, 1986; 52 FR 2684, Jan. 26, 1987; 54 FR 12189, Mar. 24, 1989; 66 FR 14074, Mar. 9, 2001]

**§ 558.254 Famphur.**

(a) *Approvals.* Type A medicated articles: 13.2 and 33.3 percent to 000061 in § 510.600(c) of this chapter.

(b) *Special considerations.* Famphur is a cholinesterase inhibitor. Do not use this product in animals simultaneously or within a few days before or after