

Amprolium and ethopabate in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(v) Amprolium 136.2 (0.015%) and ethopabate 3.6 (0.0004%).	Bacitracin 10	Broiler chickens; as an aid in the prevention of coccidiosis; growth promotion and feed efficiency.	Feed as sole ration; use as sole source of amprolium; do not feed to laying hens; as bacitracin methylene disalicylate.	047019

(2) [Reserved]

[41 FR 10990, Mar. 15, 1976]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.58, see the List of CFR Sections Affected, which appears in the Finding Aids section of the printed volume and on GPO Access.

§ 558.59 Apramycin.

(a) *Approvals.* Type A articles to sponsors identified in § 510.600(c) of this chapter as follows:

(1) 000986 for 75 grams apramycin (as apramycin sulfate) per pound for use as in paragraph (d)(1) of this section.

(2) [Reserved]

(b) [Reserved]

(c) *Related tolerances.* See § 556.52 of this chapter.

(d) *Conditions of use*—(1) *Swine*—(i) *Amount.* 150 grams per ton.

(ii) *Indications for use.* For control of porcine colibacillosis (weanling pig scours) caused by susceptible strains of *Escherichia coli*.

(iii) *Limitations.* Use for 14 days. Withdraw 28 days before slaughter.

(2) [Reserved]

[51 FR 9190, Mar. 18, 1986]

§ 558.60 Arsanilate sodium.

(a) *Approvals.* Type A medicated articles: 20, 50, or 100 percent to 015565 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.60 of this chapter.

(c) *Conditions of use.* (1) It is used for chickens and turkeys as follows:

(i) *Grams per ton.* 90 (0.01 percent).

(ii) *Indications for use.* For growth promotion and feed efficiency; improving pigmentation.

(iii) *Limitations.* Withdraw 5 days before slaughter; as sole source of organic arsenic.

(2) Arsanilate sodium may be used in accordance with the provisions of this section in the combinations provided as follows:

(i) Amprolium in accordance with § 558.55.

(ii) Zoalene in accordance with § 558.680.

[41 FR 10992, Mar. 15, 1976, as amended at 51 FR 7395, Mar. 3, 1986; 51 FR 33897, Sept. 24, 1986]

§ 558.62 Arsanilic acid.

(a) *Approvals.* Type A medicated articles to sponsors in § 510.600(c) of this chapter as follows:

(1) To 015565: 20, 50, and 100 percent for use as in the table in paragraph (c)(1), entry (ii), item 1; entry (ii), item 2; entry (iv); entry (vi); and entry (vii) of this section.

(2) To 015565: 20 percent for use as in paragraph (c)(1), entry (i); entry (ii), item 3 of this section.

(3) To 061133: 90 grams per pound arsanilic acid and 4.6 grams per pound erythromycin equivalents as erythromycin thiocyanate for use as in paragraph (c)(1), entry (iii); 90 grams per pound arsanilic acid and 9.25 grams per pound erythromycin equivalents as erythromycin thiocyanate for use as in paragraph (c)(1), entry (v).

(b) *Related tolerances.* See § 556.60 of this chapter.

(c) *Conditions of use.* (1) It is used as follows:

Arsanilic acid in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) 45 to 90 (0.005% to 0.01%).	Swine; for increased rate of weight gain and improved feed efficiency in growing swine..	Withdraw 5 days before slaughter; as sole source of organic arsenic.	015565

§ 558.76

21 CFR Ch. I (4-1-01 Edition)

Arsanilic acid in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(ii) 90 (0.01%)	1. Chickens; growth promotion and feed efficiency; improving pigmentation..do	015565
		2. Turkeys; growth promotion and feed efficiency; improving pigmentation.do	015565
		3. Swine; and aid in control of swine dysentery (hemorrhagic enteritis, bloody dysentery).do	015565
(iii)	Erythromycin 4.6	Chickens; growth promotion and feed efficiency; improving pigmentation..	As erythromycin thiocyanate; withdraw 5 days before slaughter; as sole source of organic arsenic..	012487
(iv)	Erythromycin 4.6 to 18.5.	Chickens; growth promotion and feed efficiency; improving pigmentation.	As erythromycin thiocyanate; withdraw 5 days before slaughter; as sole source of organic arsenic.	15565
(v)	Erythromycin 9.25 ...	Chickens; growth promotion and feed efficiency; improving pigmentation..	As erythromycin thiocyanate; withdraw 5 days before slaughter; as sole source of organic arsenic..	012487
(vi)	Erythromycin 92.5 ...	1. Chickens; as an aid in the prevention of chronic respiratory disease during periods of stress; growth promotion and feed efficiency; improving pigmentation.	As erythromycin thiocyanate; feed for 2 days before stress and 3 to 6 days after stress; withdraw 5 days before slaughter; as sole source of organic arsenic.	015565
		2. Chickens; as an aid in the prevention of infectious coryza; growth promotion and feed efficiency; improving pigmentation.	As erythromycin thiocyanate; feed for 7 to 14 days; withdraw 5 days before slaughter; as sole source of organic arsenic.	015565
(vii)	Erythromycin 185 ...	Chickens; as an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease; growth promotion and feed efficiency; improving pigmentation.	As erythromycin thiocyanate; feed for 5 to 8 days; do not use in birds producing eggs for food purposes; withdraw 5 days before slaughter; as sole source of organic arsenic.	15565

(2) Arsanilic acid may be used in accordance with the provisions of this section in the combinations provided as follows:

- (i) Amprolium in accordance with § 558.55.
- (ii) Amprolium and ethopabate in accordance with § 558.58.
- (iii) Bacitracin zinc in accordance with § 558.78.
- (iv) Bacitracin and zoalene in accordance with § 558.680.
- (v) Zualene in accordance with § 558.680.

[41 FR 10992, Mar. 15, 1976, as amended at 42 FR 18617, Apr. 8, 1977; 51 FR 7395, Mar. 3, 1986; 51 FR 33897, Sept. 24, 1986; 54 FR 18281, Apr. 28, 1989; 56 FR 19268, Apr. 26, 1991; 60 FR 39847, Aug. 4, 1995; 66 FR 14074, Mar. 9, 2001]

§ 558.76 Bacitracin methylene disalicylate.

- (a) *Approvals.* Type A medicated articles: 10, 25, 30, 40, 50, 60, or 75 grams per pound to 046573 in § 510.600(c) of this chapter.
- (b) *Special considerations.* The quantities of antibiotics are expressed in terms of the equivalent amount of antibiotic standard.
- (c) *Related tolerances.* See § 556.70 of this chapter.
- (d) *Conditions of use.* (1) It is used as follows: