

Food and Drug Administration, HHS

§ 573.780

petrolatum and technical white mineral oil shall not exceed the limits prescribed in paragraph (d) of this section.

(f) Petrolatum may contain any antioxidant permitted in food by regulations issued in accordance with section 409 of the act, in an amount not greater than that required to produce its intended effect.

§ 573.740 Odorless light petroleum hydrocarbons.

Odorless light petroleum hydrocarbons complying with §172.884(a) and (b) of this chapter may be safely used in an amount not in excess of that required as a component of insecticide formulations used in compliance with regulations issued in this part.

§ 573.750 Pichia pastoris dried yeast.

(a) Identity. The food additive Pichia pastoris dried yeast may be used in feed formulations of broiler chickens as a source of protein not to exceed 10 percent by weight of the total formulation.

(b) Specifications. The additive shall conform to the following percent-by-weight specifications:

- (1) Crude protein, not less than 60 percent.
(2) Crude fat, not less than 2 percent.
(3) Crude fiber, not more than 2 percent.
(4) Ash, not more than 13 percent.
(5) Moisture, not more than 6 percent.

(c) Use. To ensure safe use, the labeling of the additive and any feed additive supplement, concentrate, or premix prepared therefrom shall bear, in addition to other required information, the name of the additive, directions for use to provide not more than 10 percent by weight of the total ration, and the statement "Caution: Not to be used in layers or other poultry intended for breeding."

[58 FR 59170, Nov. 8, 1993]


§ 573.760 Poloxalene.

The food additive poloxalene may be safely used in accordance with the following prescribed conditions:

(a) The additive consists of polyoxypropylene-polyoxyethylene glycol non-

ionic block polymer meeting the following specifications:

- (1) Molecular weight range: 2,850-3,150.
(2) Hydroxyl number: 35.7-39.4.
(3) Cloud point (10 percent solution): 42° C-46° C.
(4) Structural formula:



(b) In feed as a surfactant for the flaking of feed grains when added to liquid grain conditioner in an amount not to exceed 1.0 percent of the conditioner. The conditioner is added to the feed at a rate of 1 quart per ton of feed.

(c) The label and labeling shall bear, in addition to the other information required by the Act:

- (1) The name of the additive.
(2) Adequate directions and warnings for use.

§ 573.780 Polyethylene.

(a) Identity. Polyethylene consists of basic polymers manufactured by the catalytic polymerization of ethylene.

(b) Specifications. (1) For the purposes of this section, polyethylene shall meet the specifications in item 2.1 of §177.1520(c) of this chapter.

(2) The polyethylene is designed in a pellet form in a configuration presenting maximum angular surface having the following dimensions in centimeters:

0.9±0.1 x 0.8±0.1 x 1.2±0.1

(c) Use. It is used as a replacement for roughage in feedlot rations for finishing slaughter cattle.

(d) Labeling. The labels and labeling shall bear in addition to the other information required by the Act:

- (1) The name of the additive "polyethylene roughage replacement."
(2) Adequate directions for use which shall provide for the administration of one-half pound of polyethylene pellets per head per day for 6 successive days. All natural roughage should be removed for a minimum of 12 hours prior to administration of polyethylene