

Cells, Anti-Human Globulin, or Blood Grouping Reagent.

(ii) Where a manufacturer submits data which the Director, Center for Biologics Evaluation and Research, finds adequate to establish that the mode of administration, the method of preparation, or the special nature of the product precludes or does not require a sterility test or that the sterility of the lot is not necessary to assure the safety, purity, and potency of the product, the Director may exempt a product from the sterility requirements of this section subject to any conditions necessary to assure the safety, purity, and potency of the product.

(5) *Number of final containers more than 20, less than 200.* If the number of final containers in the filling is more than 20 or less than 200, the sample shall be no less than 10 percent of the containers.

(6) *Number of final containers—20 or less.* If the number of final containers in a filling is 20 or less, the sample shall be two final containers, or the sample need be no more than one final container, provided (i) the bulk material met the sterility test requirements and (ii) after filling, it is demonstrated by testing a simulated sample that all surfaces to which the product was exposed were free of contaminating microorganisms. The simulated sample shall be prepared by rinsing the filling equipment with sterile 1.0 percent peptone solution, pH 7.1±0.1, which shall be discharged into a final container by the same method used for filling the final containers with the product.

(7) *Samples—large volume of product in final containers.* For Albumin (Human) and Plasma Protein Fraction (Human), when the volume of product in the final container is 50 milliliters or more, the final containers selected as the test sample may contain less than the full volume of product in the final containers of the filling from which the sample is taken: *Provided*, That the containers and closures of the sample are identical with those used for the filling to which the test applies, and the sample represents all stages of that filling.

(8) *Diagnostic biological products not intended for injection.* For diagnostic biological products not intended for in-

jection, (i) only the Fluid Thioglycollate Medium test incubated at 30 to 35 °C is required, (ii) the volume of material for the bulk test shall be no less than 2.0 milliliters, and (iii) the sample for the final container test shall be no less than three final containers if the total number filled is 100 or less, and, if greater, one additional container for each additional 50 containers or fraction thereof, but the sample need be no more than 10 containers.

(9) *Immune globulin preparations.* For immune globulin preparations, the test samples from the bulk material and from each final container need be no more than 2.0 ml.

(h) *Records.* The records related to the testing requirements of this section shall be prepared and maintained as required by §§211.167 and 211.194 of this chapter.

(Information collection requirements approved by the Office of Management and Budget under control number 0910-0139)

[38 FR 32056, Nov. 20, 1973, as amended at 41 FR 4015, Jan. 28, 1976; 41 FR 10428, Mar. 11, 1976; 44 FR 11754, Mar. 2, 1979; 49 FR 15187, Apr. 18, 1984; 49 FR 23834, June 8, 1984; 50 FR 4133, Jan. 29, 1985; 51 FR 44906, Dec. 15, 1986; 53 FR 12764, Apr. 19, 1988; 55 FR 11013, Mar. 26, 1990; 62 FR 48175, Sept. 15, 1997]

#### § 610.13 Purity.

Products shall be free of extraneous material except that which is unavoidable in the manufacturing process described in the approved biologics license application. In addition, products shall be tested as provided in paragraphs (a) and (b) of this section.

(a)(1) *Test for residual moisture.* Each lot of dried product shall be tested for residual moisture and shall meet and not exceed established limits as specified by an approved method on file in the biologics license application. The test for residual moisture may be exempted by the Director, Center for Biologics Evaluation and Research, when deemed not necessary for the continued safety, purity, and potency of the product.

(2) *Records.* Appropriate records for residual moisture under paragraph (a)(1) of this section shall be prepared

## §610.14

and maintained as required by the applicable provisions of §§211.188 and 211.194 of this chapter.

(b) *Test for pyrogenic substances.* Each lot of final containers of any product intended for use by injection shall be tested for pyrogenic substances by intravenous injection into rabbits as provided in paragraphs (b) (1) and (2) of this section: *Provided*, That notwithstanding any other provision of Subchapter F of this chapter, the test for pyrogenic substances is not required for the following products: Products containing formed blood elements; Cryoprecipitate; Plasma; Source Plasma; Normal Horse Serum; bacterial, viral, and rickettsial vaccines and antigens; toxoids; toxins; allergenic extracts; venoms; diagnostic substances and trivalent organic arsenicals.

(1) *Test dose.* The test dose for each rabbit shall be at least 3 milliliters per kilogram of body weight of the rabbit and also shall be at least equivalent proportionately, on a body weight basis, to the maximum single human dose recommended, but need not exceed 10 milliliters per kilogram of body weight of the rabbit, except that: (i) Regardless of the human dose recommended, the test dose per kilogram of body weight of each rabbit shall be at least 1 milliliter for immune globulins derived from human blood; (ii) for Streptokinase, the test dose shall be at least equivalent proportionately, on a body weight basis, to the maximum single human dose recommended.

(2) *Test procedure, results, and interpretation; standards to be met.* The test for pyrogenic substances shall be performed according to the requirements specified in United States Pharmacopeia XX.

(3) *Retest.* If the lot fails to meet the test requirements prescribed in paragraph (b)(2) of this section, the test may be repeated once using five other rabbits. The temperature rises recorded for all eight rabbits used in testing shall be included in determining whether the requirements are met. The lot meets the requirements for absence of pyrogens if not more than three of the eight rabbits show individual rises in temperature of 0.6 °C or more, and if the sum of the eight individual max-

## 21 CFR Ch. I (4-1-01 Edition)

imum temperature rises does not exceed 3.7 °C.

(Information collection requirements were approved by the Office of Management and Budget (OMB) and assigned OMB control number 0910-0139)

[38 FR 32056, Nov. 20, 1973, as amended at 40 FR 29710, July 15, 1975; 41 FR 10429, Mar. 11, 1976; 41 FR 41424, Sept. 22, 1976; 44 FR 40289, July 10, 1979; 46 FR 62845, Dec. 29, 1981; 49 FR 15187, Apr. 18, 1984; 50 FR 4134, Jan. 29, 1985; 55 FR 28381, July 11, 1990; 64 FR 56453, Oct. 20, 1999]

### §610.14 Identity.

The contents of a final container of each filling of each lot shall be tested for identity after all labeling operations shall have been completed. The identity test shall be specific for each product in a manner that will adequately identify it as the product designated on final container and package labels and circulars, and distinguish it from any other product being processed in the same laboratory. Identity may be established either through the physical or chemical characteristics of the product, inspection by macroscopic or microscopic methods, specific cultural tests, or in vitro or in vivo immunological tests.

### §610.15 Constituent materials.

(a) *Ingredients, preservatives, diluents, adjuvants.* All ingredients used in a licensed product, and any diluent provided as an aid in the administration of the product, shall meet generally accepted standards of purity and quality. Any preservative used shall be sufficiently nontoxic so that the amount present in the recommended dose of the product will not be toxic to the recipient, and in the combination used it shall not denature the specific substances in the product to result in a decrease below the minimum acceptable potency within the dating period when stored at the recommended temperature. Products in multiple-dose containers shall contain a preservative, except that a preservative need not be added to Yellow Fever Vaccine; Poliovirus Vaccine Live Oral; viral vaccines labeled for use with the jet injector; dried vaccines when the accompanying diluent contains a preservative; or to an Allergenic Product in 50 percent or