

§ 51.587

below the first node provided the stalk is of the length specified.

§ 51.587 Diameter.

Diameter means the greatest dimension of the stalk measured at a point 2 inches above the point of attachment of the lowest outer branch to the base.

§ 51.588 Length of seedstem.

Length of seedstem means the distance from the base of the outer branches of the stalk to the top of the actual seedstem, exclusive of any leaves or leafstems attached to the top of the seedstem.

Subpart—United States Consumer Standards for Celery Stalks

SOURCE: 18 FR 7089, Nov. 11, 1953, unless otherwise noted. Redesignated at 42 FR 32514, June 27, 1977, and further redesignated at 46 FR 63203, Dec. 31, 1981.

GENERAL

§ 51.595 General.

These standards do not apply to celery hearts.

GRADES

§ 51.596 U.S. Grade AA.

U.S. Grade AA shall consist of stalks of celery of similar varietal characteristics, which are well developed, and have good heart formation; which are clean, well trimmed, fairly compact, and are free from blackheart, brown stem, decay (except dry type crater rot), doubles, and from damage caused by crater rot, wilting, cutworms, freezing, suckers, growth cracks, hollow crown, pithy branches, seedstems, rust, cracked stem, other diseases, insects or mechanical or other means.

(a) The average midrib length of the outer whorl of branches on stalks in this grade shall be not less than 7 inches. (See §§ 51.599 and 51.600.)

(b) Incident to proper grading and handling other than for average midrib length of branches not more than 5 percent, by count, of the stalks in any lot may fail to meet the requirements of the grade including not more than 1 percent for stalks affected by moist type decay. In addition, not more than

7 CFR Ch. I (1-1-01 Edition)

3 percent, by count, of the stalks in any lot may fail to meet the requirements as to average midrib length of the stalks.

§ 51.597 U.S. Grade A.

U.S. Grade A shall consist of stalks of celery of similar varietal characteristics which are fairly well developed and have fairly good heart formation; which are clean, well trimmed, and not badly spread, and which are free from blackheart, decay (except dry type crater rot), doubles, and from damage caused by crater rot, brown stem, wilting, cutworms, freezing, suckers, growth cracks, hollow crown, pithy branches, seedstems, rust, cracked stem, other diseases, insects, or mechanical or other means.

(a) The average midrib length of the outer whorl of branches on stalks in this grade shall be not less than 5 inches. (See §§ 51.599 and 51.600.)

(b) Incident to proper grading and handling other than for average midrib length of branches, not more than 5 percent, by count, of the stalks in any lot may fail to meet the requirements of the grade including not more than 1 percent for stalks affected by moist type decay. In addition, not more than 3 percent, by count, of the stalks in any lot may fail to meet the requirements as to average midrib length of the stalks.

§ 51.598 U.S. Grade B.

U.S. Grade B shall consist of stalks of celery of similar varietal characteristics which are fairly well developed, which are clean, well trimmed and free from blackheart, decay (except dry type crater rot), doubles, and from serious damage caused by crater rot, brown stem, wilting, cutworms, freezing, suckers, growth cracks, hollow crown, pithy branches, seedstems, rust, cracked stem, other diseases, insects, or mechanical or other means.

(a) The average midrib length of the outer whorl of branches on stalks in this grade shall be not less than 4 inches. (See §§ 51.599 and 51.600.)

(b) Incident to proper grading and handling other than for average midrib length of branches, not more than 5 percent, by count, of the stalks in any lot may fail to meet the requirements

Agricultural Marketing Service, USDA

§ 51.609

of this grade including not more than 1 percent for stalks affected by moist type decay. In addition not more than 3 percent, by count, of the stalks in any lot may fail to meet the requirements as to average midrib length of the stalks.

en Self Blanching types must not be mixed.

BLANCHING; LENGTH OF STALKS

§ 51.604 Well developed.

§ 51.599 Blanching.

Well developed means that the outer branches are of good width in relation to the length of midribs and type of celery.

There are no requirements in the grades as to blanching. However, celery stalks may be classed as “green” when they have a medium to dark green appearance, fairly “well blanched” when the midrib portions of the branches on the stalks are generally of a light greenish to creamy white color, or “well blanched” when the midrib portions of the branches on the stalks are generally of a creamy white color. Not more than 5 percent of the stalks in any lot may fail to meet the requirements of any of the above classes.

§ 51.605 Good heart formation.

Good heart formation means that the stalk has a reasonable number of stocky inner heart branches for its size.

§ 51.600 Length of stalks.

§ 51.606 Clean.

Clean means that the stalk is practically free from dirt or other foreign materials. Stalks shall be permitted to have a small amount of dirt on the inside of the branches or in the heart branches which cannot be removed by good commercial methods of washing.

There are no requirements in the grades as to stalk length. However, when the stalk length is specified it shall be determined by measuring the distance from where the main root is cut off, to a point which represents the average length of the longest branches and leaves expressed in terms of the nearest whole inch. Incident to proper sizing, not more than 5 percent, by count, of the stalks in any lot may fail to meet any specified stalk length.

§ 51.607 Well trimmed.

Well trimmed means that the outside coarse and damaged branches have been removed and that the root or roots have been neatly trimmed to a reasonable length for the size of the stalk.

OFF-GRADE

§ 51.608 Fairly compact.

Fairly compact means that the branches are fairly close together on the stalk.

§ 51.601 Off-Grade celery.

Celery stalks which fail to meet the requirements of any of the foregoing grades shall be Off-Grade celery stalks.

§ 51.609 Damage.

Damage means any injury or defect which materially affects the appearance, or edible or shipping quality. Any one of the following defects, or any combination of defects, the seriousness of which exceeds the maximum allowed for any one defect shall be considered as damage:

DEFINITIONS

(a) Crater rot, when moist, or when occurring on more than 2 branches, or when aggregating more than two-thirds of a square inch on the branch or branches.

§ 51.602 Stalk.

Stalk means an individual plant.

(b) Cutworms, when the worms are present, or when worm injury occurs on the heart branches, or when occurring on the midrib portion of more than two branches, or when aggregating more than one-half of a square inch on the midrib portion of the branch or branches.

§ 51.603 Similar varietal characteristics.

Similar varietal characteristics means that the stalks in any container have the same character of growth. For example, celery of Giant Pascal and Gold-