

§ 520.1484

initial intravenous dosage is 5 milligrams per kilogram of body weight.

(ii) For oral dosage only, administer 10 milligrams naproxen per kilogram of animal body weight twice daily as a top dressing in the animal's feed for up to 14 consecutive days.

(3) Not for use in horses intended for food.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[41 FR 14188, Apr. 2, 1976, as amended at 51 FR 24525, July 7, 1986; 61 FR 5506, Feb. 13, 1996]

§ 520.1484 Neomycin sulfate soluble powder.

(a) *Specifications.* Neomycin sulfate soluble powder contains 20.3 grams of neomycin sulfate (equivalent to 14.2 grams of neomycin base) per ounce.

(b) *Sponsors.* See 000069, 046573, 051259, and 061133 in § 510.600(c) of this chapter for use as in paragraph (d)(1) of this section. See 000009 for use as in paragraphs (d)(1) and (d)(2) of this section.

(c) *Related tolerances.* See § 556.430 of this chapter.

(d) *Conditions of use*—(1) *Cattle (excluding veal calves), swine, sheep, and goats.*

(i) *Amount.* 10 milligrams of neomycin sulfate per pound of body weight per day (22 milligrams per kilogram) in divided doses for a maximum of 14 days.

(ii) *Indications for use.* For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves), swine, sheep, and goats.

(iii) *Limitations.* Add to drinking water or milk; not for use in liquid supplements. Prepare a fresh solution daily. If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 14 consecutive days. Discontinue treatment prior to slaughter as follows: Cattle (not for use in veal calves), 1 day; sheep, 2 days; swine and goats, 3 days.

(2) *Turkeys*—(i) *Amount.* 10 milligrams of neomycin sulfate per pound of body

21 CFR Ch. I (4–1–02 Edition)

weight per day (22 milligrams per kilogram) for 5 days.

(ii) *Indications for use.* For the control of mortality associated with *E. coli* organisms susceptible to neomycin sulfate in growing turkeys.

(iii) *Limitations.* Add to drinking water; not for use in liquid supplements. Prepare a fresh solution daily. If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 5 consecutive days.

[64 FR 31497, June 11, 1999, as amended at 66 FR 14073, Mar. 9, 2001]

§ 520.1485 Neomycin sulfate oral solution.

(a) *Specifications.* Each milliliter contains 200 milligrams of neomycin sulfate (equivalent to 140 milligrams of neomycin base).

(b) *Sponsors.* See Nos. 000009, 051259, and 059130 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.430 of this chapter.

(d) *Conditions of use*—(1) *Amount.* 10 milligrams of neomycin sulfate per pound of body weight per day in divided doses for a maximum of 14 days.

(2) *Indications for use.* For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin in cattle (excluding veal calves), swine, sheep, and goats.

(3) *Limitations.* Administer undiluted or in drinking water. Prepare a fresh solution daily. If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 14 consecutive days. Discontinue treatment prior to slaughter as follows: 1 day for cattle, 2 days for sheep, and 3 days for swine and goats.

[58 FR 38972, July 21, 1993, as amended at 60 FR 3079, Jan. 13, 1995; 61 FR 31398, June 20, 1996; 62 FR 60657, Nov. 12, 1997; 63 FR 45944, Aug. 28, 1998; 65 FR 45877, July 26, 2000; 65 FR 53581, Sept. 5, 2000]