§520.1855 Ponazuril.

(a) Specifications. Each gram of paste contains 150 milligrams (mg) ponazuril.

(b) Sponsor. See No. 000859 in §510.600(c) of this chapter.

(c) Conditions of use in horses—(1) Amount. 5 mg per kilogram body weight, daily for 28 days.

(2) Indications for use. For the treatment of equine protozoal myeloencephalitis caused by Sarcocystis neurona.

(3) *Limitations*. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[66 FR 43774, Aug. 21, 2001]

§ 520.1870 Praziquantel tablets.

(a) Specifications. Each dog tablet contains 34 milligrams (mg) of praziquantel; each cat tablet contains 11.5 or 23 mg of praziquantel.

(b) Sponsor. See No. 000859 in §510.600(c) of this chapter.

(c) Conditions of use—(1) Dogs—(i) Indications for use. For removal of canine cestodes Dipylidium caninum and Taenia pisiformis. If labeled for use by or on the order of a licensed veterinarian, for removal of the canine cestode Echinococcus granulosus, and for removal and control of the canine cestode Echinococcus multilocularis.

(ii) Dosage. Dogs 5 pounds and under, $\frac{1}{2}$ tablet (17 mg); 6 to 10 pounds, 1 tablet (34 mg); 11 to 15 pounds, 1 $\frac{1}{2}$ tablets (51 mg); 16 to 30 pounds, 2 tablets (68 mg); 31 to 45 pounds, 3 tablets (102 mg); 46 to 60 pounds, 4 tablets (136 mg); over 60 pounds, 5 tablets maximum (170 mg).

(iii) Limitations. Administer directly by mouth or crumbled and in feed. Not intended for use in puppies less than 4 weeks of age. For over-the-counter (OTC) use: Consult your veterinarian before administering tablets to weak or debilitated animals, and for assistance in the diagnosis, treatment, and control of parasitism. For prescription use: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) Cats—(i) Indications for use. For removal of feline cestodes Dipylidium caninum and Taenia taeniaeformis. 21 CFR Ch. I (4–1–02 Edition)

(ii) *Dosage*. Cats 4 pounds and under, 11.5 mg; 5 to 11 pounds, 23 mg; over 11 pounds, 34.5 mg.

(iii) *Limitations*. Administer directly by mouth or crumbled and in feed. Not intended for use in kittens less than 6 weeks of age. For OTC use: Consult your veterinarian before administering tablets to weak or debilitated animals, and for assistance in the diagnosis, treatment, and control of parasitism.

[46 FR 60570, Dec. 11, 1981, as amended at 47
FR 26377, June 18, 1982; 55 FR 2234, Jan. 23, 1990; 58 FR 7864, Feb. 10, 1993; 58 FR 42853, Aug. 12, 1993]

§ 520.1871 Praziquantel/pyrantel pamoate tablets.

(a) *Specifications*. Each cat tablet contains 18.2 milligrams (mg) praziquantel with 72.6 mg pyrantel (as pyrantel pamoate).

(b) Sponsor. See 000859 in 510.600(c) of this chapter.

(c) Conditions of use—(1) Cats—(i) Dosage. 1.5 to 1.9 pounds, 1/4 tablet; 2 to 3 pounds, 1/2 tablet; 4 to 8 pounds, 1 tablet; 9 to 12 pounds, 1 1/2 tablets; 13 to 16 pounds, 2 tablets.

(ii) Indications for use. For removal of tapeworms (Dipylidium caninum and Taenia taeniaeformis), hookworms (Ancylostoma tubaeforme), and large roundworms (Toxocara cati) in cats and kittens.

(iii) Limitations. Not for use in kittens less than 1 month of age or weighing less than 1.5 pounds. May be given directly by mouth or in a small amount of food. Do not withhold food prior to or after treatment. If reinfection occurs, treatment may be repeated. Consult your veterinarian before giving to sick or pregnant animals. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(2) [Reserved]

[58 FR 58652, Nov. 3, 1993]

§ 520.1872 Praziquantel, pyrantel pamoate, and febantel tablets.

(a) *Specifications*. Each tablet contains either:

(1) Tablet No. 1: 22.7 milligrams praziquantel, 22.7 milligrams pyrantel base, and 113.4 milligrams febantel; or