

**§ 520.1920**

**21 CFR Ch. I (4-1-02 Edition)**

clinically recognizable lesion in certain entities in dogs.<sup>1</sup>

(3) *Limitations.* The tablets may be administered whole or crushed and mixed with the food. When convulsions are frequent, the dosage should be divided and administered at intervals. Reduction in dosage should be made gradually and never be abruptly discontinued. Do not use in feline species, as primidone appears to have a specific neurotoxicity in cats. Federal law restricts this drug to use by or on the order of a licensed veterinarian.<sup>1</sup>

[42 FR 61594, Dec. 6, 1977, as amended at 43 FR 55386, Nov 28, 1978; 46 FR 8467, Jan. 27, 1981; 46 FR 57477, Nov. 24, 1981; 53 FR 40727, Oct. 18, 1988; 56 FR 37473, Aug. 7, 1991; 62 FR 35076, June 30, 1997]

**§ 520.1920 Prochlorperazine, isopropamide sustained release capsules.**

(a) *Specifications.* Prochlorperazine, isopropamide sustained release capsules contain either:

(1) 3.33 milligrams of prochlorperazine (as the dimaleate) and 1.67 milligrams of isopropamide (as the iodide), or

(2) 10 milligrams of prochlorperazine (as the dimaleate) and 5 milligrams of isopropamide (as the iodide).

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used for the treatment of dogs in which gastrointestinal disturbances are associated with emotional stress.

(2)(i) Capsules described in paragraph (a)(1) of this section are administered orally to dogs weighing from 4 to 15 pounds at the rate of 1 capsule twice daily. These capsules are administered orally to dogs weighing from 16 to 30 pounds at the rate of 1 or 2 capsules twice daily. For dogs weighing less than 4 pounds, administer orally an appropriate fraction of the contents of one of these capsules.

(ii) Capsules described in paragraph (a)(2) of this section are given to dogs weighing 30 pounds and over at the rate of 1 capsule twice daily.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 56 FR 50653, Oct. 8, 1991; 60 FR 55659, Nov. 2, 1995]

**§ 520.1921 Prochlorperazine, isopropamide, with neomycin sustained-release capsules.**

(a) *Specifications.* Each capsule contains either:

(1) Capsule No. 1: 3.33 milligrams of prochlorperazine (as the dimaleate), 1.67 milligrams of isopropamide (as the iodide), and 25 milligrams of neomycin base (as the sulfate); or

(2) Capsule No. 3: 10 milligrams of prochlorperazine (as the dimaleate), 5 milligrams of isopropamide (as the iodide), and 75 milligrams of neomycin base (as the sulfate).

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use—*(1) *Amount.* Administer capsules orally twice daily to dogs as follows:

Animal weight (pounds)	Number of capsules per dose	
	Capsule No. 1	Capsule No. 3
10 to 20 .....	1	
20 to 30 .....	2	
Over 30 .....	3	1
Over 60 .....		2

(2) *Indications for use.* For treatment of dogs in which infectious bacterial gastroenteritis is associated with emotional stress.

(3) *Limitations.* Do not continue medication longer than 5 days. Overdosage or prolonged administration may produce nephrotoxicity as manifested by albuminuria, presence of granular casts and depressed urinary output. If it is desirable to administer a vasoconstrictor, norepinephrine is the drug of choice. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[49 FR 14103, Apr. 10, 1984, as amended at 56 FR 50653, Oct. 8, 1991; 60 FR 55659, Nov. 2, 1995]

**§ 520.1962 Promazine hydrochloride.**

(a)(1) *Chemical name.* 10-[3-(Dimethylamino)propyl]phenothiazine monohydrochloride.

(2) *Specifications.* Conforms to N.F. XII.

(3) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(4) [Reserved]

(5) *Conditions of use.* (i) The drug is used for quieting excitable, unruly, or