

**§ 520.2158b**

may develop a tolerance for streptomycin. Consult a veterinarian or animal pathologist for diagnosis.

(2) *Chickens*—(i) *Amount*. 10 to 15 mg/pound of body weight (0.6 to 0.9 grams per gallon).

(ii) *Indications for use*. Treatment of nonspecific infectious enteritis caused by organisms susceptible to streptomycin.

(iii) *Limitations*. Chickens: Do not administer for more than 5 days. Withdraw 4 days before slaughter. Do not administer to chickens producing eggs for human consumption. Prepare fresh solution daily. As sole source of streptomycin. Warning: Certain strains of bacteria may develop a tolerance for streptomycin. Consult a veterinarian or animal pathologist for diagnosis.

[57 FR 37327, Aug. 18, 1992, as amended at 58 FR 47211, Sept. 8, 1993; 63 FR 51821, Sept. 29, 1998]

**§ 520.2158b Dihydrostreptomycin tablets.**

(a) *Specifications*. Each tablet contains 37.5 milligrams dihydrostreptomycin (as the sulfate) with 375 milligrams chlorhexidine dihydrochloride.

(b) *Sponsor*. See No. 000856 in § 510.600(c) of this chapter.

(c) *Related tolerances*. See §§ 556.120 and 556.200 of this chapter.

(d) *Conditions of use. Calves*—(1) *Amount*. 150 milligrams of dihydrostreptomycin and 1.5 grams of chlorhexidine dihydrochloride per 100 pounds of body weight per day.

(2) *Indications for use*. Treatment of bacterial scours in calves.

(3) *Limitations*. Administer orally once a day for 5 days; withdraw 3 days before slaughter.

[57 FR 37327, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992]

**§ 520.2158c Dihydrostreptomycin oral suspension.**

(a) *Specifications*. Each milliliter contains 1.25 milligrams dihydrostreptomycin (as the sulfate) with 12.5 milligrams chlorhexidine dihydrochloride.

(b) *Sponsor*. See No. 000856 in § 510.600(c) of this chapter.

(c) *Related tolerances*. See §§ 556.120 and 556.200 of this chapter.

(d) *Conditions of use. Calves*—(1) *Amount*. 150 milligrams of dihydro-

**21 CFR Ch. I (4–1–02 Edition)**

streptomycin and 1.5 grams of chlorhexidine dihydrochloride per 100 pounds of body weight per day.

(2) *Indications for use*. Treatment of bacterial scours in calves.

(3) *Limitations*. Administer orally once a day for 5 days; withdraw 3 days before slaughter.

[57 FR 37327, Aug. 18, 1992]

**§ 520.2160 Styrylpyridinium, diethylcarbazine oral dosage forms.**

**§ 520.2170 Sulfabromomethazine sodium boluses.**

(a) *Specifications*. Each bolus contains 15 grams of sulfabromomethazine sodium.

(b) *Related tolerance*. See § 556.620 of this chapter.

(c) *Sponsor*. See No. 050604 in § 510.600(c) of this chapter.

(d) *NAS/NRC status*. These conditions of use are NAS/NRC reviewed and found effective. NADA's for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(e) *Conditions of use. Cattle*—(1) *Amount*. 90 milligrams per pound body weight.

(2) *Indications for use*. Treatment of necrotic pododermatitis (foot rot) and calf diphtheria caused by *Fusobacterium necrophorum*; colibacillosis (scours) caused by *Escherichia coli*; bacterial pneumonia and bovine respiratory disease complex (shipping fever complex) associated with *Pasteurella* spp.; acute metritis and acute mastitis caused by *Streptococcus* spp.

(3) *Limitations*. Administer orally; repeat in 48 hours if necessary; milk taken from animals within 96 hours (8 milkings) of latest treatment must not be used for food; do not administer within 18 days of slaughter; discontinue use if hematuria, crystalluria or severe depression are noticed; if signs persist after 2 or 3 days consult a veterinarian.

[47 FR 30243, July 13, 1982, as amended at 62 FR 63270, Nov. 28, 1997]