§ 520.2345

- (c) Conditions of use—(1) Amount. Administer one tablet orally per 4 pounds of body weight.¹
- (2) *Indications for use.* Use in dogs and cats as an aid in treatment of bacterial pneumonia and bacterial enteritis when caused by organisms sensitive to sulfisoxazole.¹
- (3) Limitations. Repeat dosage at 24-hour intervals until 2 to 3 days after disappearance of clinical symptoms. (Administration of one-half daily dosage at 12-hour intervals or one-third daily dosage at 8-hour intervals will provide a more constant blood level.) Provide adequate supply of drinking water. If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian.¹

[43 FR 60895, Dec. 29, 1978]

§ 520.2345 Tetracycline oral dosage forms.

§ 520.2345a Tetracycline hydrochloride capsules.

- (a) *Specifications*. Each capsule contains 50, 100, 125, 250, or 500 milligrams of tetracycline hydrochloride.
- (b) Sponsor. See §510.600(c) of this chapter for identification of the sponsors:
- (1) To No. 000009: 250 milligrams per capsule.
- (2) To No. 000069: 125, 250, and 500 milligrams per capsule.
- (3) To No. 000115: 50, 100, 250, and 500 milligrams per capsule.
- (c) Conditions of use. Dogs—(1) Amount. 25 milligrams per pound of body weight per day in divided doses every 6 hours.
- (2) Indications for use. Treatment of infections caused by organisms sensitive to tetracycline hydrochloride, such as bacterial gastroenteritis due to *E. coli* and urinary tract infections due to *Staphylococcus* spp. and *E. coli*.
- (3) Limitations. Administer orally; continue treatment until symptoms of the disease have subsided and the temperature is normal for 48 hours; not for use in animals raised for food produc-

tion; Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37327, Aug. 18, 1992, as amended at 59 FR 59365, Nov. 17, 1994; 63 FR 5255, Feb. 2, 1998]

$\S 520.2345b$ Tetracycline tablets.

- (a) *Specifications*. Each tablet contains 100, 250, or 500 milligrams of tetracycline (as the hydrochloride).
- (b) *Sponsor*. For 100, 250, or 500 milligrams per tablet, see No. 000069 in §510.600(c) of this chapter. For 250 milligrams per tablet, see No. 000009 in §510.600(c) of this chapter.
- (c) Conditions of use. Dogs—(1) Amount. 25 milligrams per pound of body weight per day in divided doses every 6 hours.
- (2) Indications for use. Treatment of infections caused by organisms sensitive to tetracycline hydrochloride, such as bacterial gastroenteritis due to *E. coli* and urinary tract infections due to *Staphylococcus* spp. and *E. coli*.
- (3) Limitations. Administer orally; continue treatment until symptoms of the disease have subsided and temperature is normal for 48 hours; not for use in animals raised for food production; Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37327, Aug. 18, 1992]

§520.2345c Tetracycline boluses.

- (a) *Specifications*. Each bolus contains 500 milligrams of tetracycline (as the hydrochloride).
- (b) *Sponsors*. See No. 010042 in §510.600(c) of this chapter for use as in paragraph (d)(1) of this section. See No. 000009 in §510.600(c) of this chapter for use as in paragraph (d)(2) of this section.
- (c) Related tolerances. See §556.720 of this chapter.
- (d) Conditions of use. Calves—(1) Amount. 10 milligrams per pound of body weight per day in divided doses.
- (i) Indications for use. Control and treatment of bacterial enteritis (scours) caused by *E. coli* and bacterial pneumonia caused by *Pasteurella* spp., *Hemophilus* spp., and *Klebsiella* spp.
- (ii) *Limitations*. Administer orally for 3 to 5 days; do not slaughter animals

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by §514.111 of this chapter, but may require bioequivalency and safety information.