

(b) *Sponsor*. See No. 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use*. (1) It is administered to horses by stomach tube or as a drench at the rate of 1 fluid ounce of suspension per 100 pounds of body weight for the control of large strongyles, small strongyles, pinworms, *Strongyloides* and ascarids (including members of the genera *Strongylus spp.*, *Cyathostomum spp.*, *Cylicobrachytus spp.* and related genera *Craterostomum spp.*, *Oesophagodontus spp.*, *Poteriostomum spp.*, *Oxyuris spp.*, *Strongyloides spp.*, and *Parascaris spp.*).

(2) Do not use in horses intended to be used for food purposes.

(3) For use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 62 FR 63271, Nov. 28, 1997]

§ 520.2380e Thiabendazole with trichlorfon.

(a) *Specifications*. The drug contains 5 grams of thiabendazole with 4.5 grams of trichlorfon, or 20 grams of thiabendazole with 18 grams of trichlorfon.

(b) *Sponsor*. See No. 017135 in § 510.600(c) of this chapter.

(c) *Conditions of use*. (1) Used for the treatment and control of bots (*Gasterophilus spp.*), large strongyles (*Strongylus spp.*), small strongyles (genera *Cyathostomum*, *Cylicobrachytus*, *Craterostomum*, *Oesophagodontus*, *Poteriostomum*), pinworms (*Oxyuris spp.*, *Strongyloides spp.*), and ascarids (*Parascaris spp.*) in horses.

(2) Administer 2 grams of thiabendazole with 1.8 grams of trichlorfon per 100 pounds of body weight sprinkled on the animals' usual daily ration of feed, or may be mixed in 5 to 10 fluid ounces of water and administered by stomach tube or drench.

(3) Do not re-treat more than once every 30 days, preferably every 6 to 8 weeks.

(4) Do not treat animals if sick or debilitated; less than 4 months of age; or mares in last month of pregnancy.

(5) Do not administer intravenous anesthetics, especially muscle relaxants, within 2 weeks of use.

(6) Not for animals intended for food use.

(7) Do not use within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(8) If the label bears directions for administration of the drug by stomach tube or drench it shall also bear the statement: Caution; Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 23071, May 28, 1975, as amended at 48 FR 48229, Oct. 18, 1983]

§ 520.2380f Thiabendazole, piperazine phosphate powder.

(a) *Specifications*. Each ounce of water dispersible powder contains 6.67 grams of thiabendazole and 8.33 grams of piperazine (as piperazine phosphate).

(b) *Sponsor*. See No. 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*. 2 grams of thiabendazole and 2.5 grams of piperazine (0.3 ounce of powder) per 100 pounds of body weight.

(2) *Indications for use*. Treatment of infections of large strongyles (genus *Strongylus*), small strongyles (genera *Cyathostomum*, *Cylicobrachytus*, and related genera *Craterostomum*, *Oesophagodontus*, *Poteriostomum*), pinworms (*Oxyuris*), threadworms (*Strongyloides*), and ascarids (*Parascaris*) in horses.

(3) *Limitations*. Use a single oral dose. Administer as a drench or by stomach tube suspended in 1 pint of warm water; by dose syringe suspended in ½ ounce of water for each 100 pounds of body weight; or sprinkled over a small amount of daily feed. Not for animals intended for food use. If the label bears directions for administration by stomach tube or drench, it shall also bear the statement "Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian;" if not labeled for use by stomach tube or drench, the label shall bear the statement, "Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism."

[46 FR 18963, Mar. 27, 1981, as amended at 46 FR 52330, Oct. 27, 1981; 62 FR 63271, Nov. 28, 1997]

§ 520.2455 Tiamulin soluble powder.

(a) *Specifications*. A water-soluble powder containing 45 percent tiamulin