

§ 520.2520

(3) The labeling shall comply with the requirements of §510.410 of this chapter.

(4) Not for use in horses intended for food.

(5) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[41 FR 24884, June 21, 1976, as amended at 50 FR 41489, Oct. 11, 1985; 51 FR 26002, July 18, 1986]

§ 520.2520 Trichlorfon oral dosage forms.

§ 520.2520b Trichlorfon and atropine.

(a) *Chemical name.* (1) For trichlorfon: *O,O*-Dimethyl 2,2,2-trichloro-1-hydroxyethyl phosphonate.

(2) For atropine: Atropine N.F.

(b) *Sponsor.* See No. 000856 in §510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used for the treatment of *Syphacia obvelata* (pinworm) in laboratory mice.

(2) It is administered in distilled water as sole source of drinking water continuously for 7 to 14 days at 1.67 grams of trichlorfon and 7.7 milligrams of atropine per liter.

(3) Prepare fresh solution every 3 days. Do not use simultaneously with other drugs, insecticides, pesticides, or chemicals having cholinesterase activity, nor within 7 days before or after treatment with any other cholinesterase inhibitor.

(4) Restricted to use by or on the order of a licensed veterinarian.

§ 520.2520e Trichlorfon boluses.

(a) *Specifications.* Each bolus contains either 7.3, 10.9, 14.6, or 18.2 g of trichlorfon.

(b) *Sponsor.* See 000856 in §510.600(c) of this chapter.

(c) *Special considerations.* Trichlorfon is a cholinesterase inhibitor. Do not use this product on animals simultaneously with, or within 2 weeks, before or after treatment with or exposure to, neuromuscular depolarizing agents (i.e., succinylcholine) or to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(d) *NAS/NRC status.* Use of this drug has been NAS/NRC reviewed and found effective. Applications for these uses

21 CFR Ch. I (4-1-02 Edition)

need not include effectiveness data as specified by §514.111 of this chapter.

(e) *Conditions of use*—(1) *Amount.* 18.2 milligrams per pound of body weight, except for strongyles use 36.4 milligrams per pound of body weight.

(2) *Indications for use.* For horses for removal of bots (*Gastrophilus nasalis*, *Gastrophilus intestinalis*), large strongyles (*Strongylus vulgaris*), small strongyles, large roundworms (ascarids, *Parascaris equorum*), and pinworms (*Oxyuris equi*).

(3) *Limitations.* Do not fast horses before or after treatment. Treatment of mares in late pregnancy is not recommended. Surgery or any severe stress should be avoided for at least 2 weeks before or after treatment. Do not administer to sick, toxic, or debilitated horses. Not to be used in horses intended for use as food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[45 FR 48127, July 18, 1980]

§ 520.2520f Trichlorfon granules.

(a) *Specifications.* Each package contains either 18.2 or 36.4 g of trichlorfon.

(b) *Sponsor.* See 000856 in §510.600(c) of this chapter.

(c) *Special considerations.* Trichlorfon is a cholinesterase inhibitor. Do not use this product on animals simultaneously with, or within 2 weeks before or after treatment with or exposure to neuromuscular depolarizing agents (i.e., succinylcholine) or to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(d) *NAS/NRC status.* Use of this drug has been NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by §514.111 of this chapter.

(e) *Conditions of use*—(1) *Amount.* 18.2 milligrams per pound of body weight.

(2) *Indications for use.* For horses for removal of bots (*Gastrophilus nasalis*, *Gastrophilus intestinalis*), large roundworms (ascarids, *Parascaris equorum*), and pinworms (*Oxyuris equi*).

(3) *Limitations.* Do not fast horses before or after treatment. Treatment of mares in late pregnancy is not recommended. Surgery or any severe stress should be avoided for at least 2 weeks before or after treatment. Do