## §520.88d

(d) Conditions of use. Swine—(1) Amount. 40 milligrams orally, twice a day using a dosing pump.

(2) *Indications for use*. Treatment of baby pigs under 10 pounds for porcine colibacillosis caused by *Escherichia coli* susceptible to amoxicillin.

(3) Limitations. Treat animals for 48 hours after all symptoms have subsided but not beyond 5 days. Do not slaughter during treatment or for 15 days after latest treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37319, Aug. 18, 1992, as amended at 60 FR 55658, Nov. 2, 1995]

## § 520.88d Amoxicillin trihydrate soluble powder.

(a) *Specifications*. Each gram contains amoxicillin trihydrate equivalent to 115.4 milligrams of amoxicillin.

(b) Sponsor. See No. 000069 in §510.600(c) of this chapter.

(c) *Related tolerances*. See §556.38 of this chapter.

(d) Conditions of use. Preruminating calves including veal calves—(1) Amount.
400 milligrams per 100 pounds of body weight twice daily.

(2) Indications for use. Treatment of bacterial enteritis when due to susceptible *Escherichia coli* in preruminating calves including veal calves.

(3) Limitations. Administer by drench or by mixing in milk. Treatment should be continued for 48 hours after all symptoms have subsided but not to exceed 5 davs. For 1150 in preruminating calves including veal calves only, not for use in other animals which are raised for food production. Do not slaughter animals during treatment or for 20 days after the latest treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37319, Aug. 18, 1992; 57 FR 42623, Sept.
15, 1992; 58 FR 18304, Apr. 8, 1993, as amended at 60 FR 55658, Nov. 2, 1995; 62 FR 5525, Feb. 6, 1997]

## § 520.88e Amoxicillin trihydrate boluses.

(a) *Specifications*. Each bolus contains the equivalent of 400 milligrams of amoxicillin.

21 CFR Ch. I (4–1–02 Edition)

(b) *Sponsor*. See No. 000069 in §510.600(c) of this chapter.

(c) *Related tolerances*. See §556.38 of this chapter.

(d) Conditions of use. Preruminating calves including veal calves—(1) Amount.
400 milligrams per 100 pounds of body weight twice daily.

(2) Indications for use. Treatment of bacterial enteritis when due to susceptible *Escherichia coli* in preruminating calves including veal calves.

(3) Limitations. For oral use in preruminating calves including veal calves only, not for use in other animals which are raised for food production. Treatment should be continued for 48 hours after all symptoms have subsided but not to exceed 5 days. Do not slaughter animals during treatment or for 20 days after the latest treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37320, Aug. 18, 1992, as amended at 60
 FR 55659, Nov. 2, 1995; 62 FR 5526, Feb. 6, 1997]

## §520.88f Amoxicillin trihydrate tablets.

(a) *Specifications*. Each tablet contains amoxicillin trihydrate equivalent to 50, 100, 200, or 400 milligrams of amoxicillin.

(b) *Sponsor*. See Nos. 000031 or 000093 in §510.600(c) of this chapter.

(c) Conditions of use—(1) Dogs—(i) Amount. 5 milligrams per pound of body weight twice a day.

(ii) Indications for use. Treatment of bacterial dermatitis due to Staphylococcus aureus, Streptococcus spp., Staphylococcus spp., and Escherichia coli; and soft tissue infections (abscesses, wounds, lacerations) due to S. aureus, Streptococcus spp., E. coli, Proteus mirabilis, and Staphylococcus spp.

(iii) *Limitations*. Use for 5 to 7 days or 48 hours after all symptoms have subsided. If no improvement is seen in 5 days, review diagnosis and change therapy. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[57 FR 37320, Aug. 18, 1992, as amended at 62 FR 13302, Mar. 20, 1997]