(c) Conditions of use—(1) Amount. (i) Dogs: Administer orally from 0.0625 to 0.25 milligram daily in divided doses.

(ii) *Cats:* Administer orally from 0.03125 to 0.125 milligram daily in divided doses.

(2) Indications for use. (i) Dogs: It is used for musculoskeletal conditions due to inflammation of muscles or joints and accessory structures, where permanent structural changes do not exist, such as arthritis, the disc syndrome, and myositis.

(ii) *Dogs and cats*: It is used in certain acute and chronic dermatoses of varying etiology to help control the pruritus, irritation, and inflammation associated with these conditions.

(3) Limitations. Do not use in viral infections. Anti-inflammatory action of corticosteroids may mask signs of infection. Do not use in animals with tuberculosis. chronic nephritis. cushingoid syndrome, or where peptic ulcers occur, except for emergency therapy. Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

 $[44\ {\rm FR}\ 7131,\ {\rm Feb.}\ 6,\ 1979,\ {\rm as}\ {\rm amended}\ {\rm at}\ 61\ {\rm FR}\ 5506,\ {\rm Feb.}\ 13,\ 1996]$ 

## § 520.970 Flunixin oral dosage forms.

## §520.970a Flunixin meglumine granules.

(a) *Specifications*. Each 10-gram packet contains flunixin meglumine equivalent to 250 milligrams of flunixin.

(b) *Sponsor*. No. 000061 in §510.600(c) of this chapter.

(c) Conditions of use—(1) Amount. 0.5 milligram of flunixin per pound of body weight (one packet per 500 pounds) per day.

(2) *Indications for use*. For alleviation of inflammation and pain associated with musculoskeletal disorders in the horse.

(3) *Limitations*. Administer daily dose for up to 5 days by sprinkling on small

21 CFR Ch. I (4–1–02 Edition)

amount of feed. The effect of this drug on pregnancy has not been determined. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[44 FR 36381, June 22, 1979. Redesignated at 50 FR 38114, Sept. 20, 1985, and amended at 52 FR 7832, Mar. 13, 1987]

## § 520.970b Flunixin meglumine paste.

(a) *Specifications*. Each 30-gram syringe contains flunixin meglumine equivalent to 1,500 milligrams of flunixin.

(b) Sponsor. No. 000061 in 510.600(c) of this chapter.

(c) Conditions of use. Horses—(1) Amount. 0.5 milligram of flunixin per pound of body weight daily.

(2) Indications for use. For alleviation of inflammation and pain associated with musculoskeletal disorders.

(3) Limitations. For oral use only. Treatment should not exceed 5 consecutive days. The effect of this drug on pregnancy has not been determined. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[50 FR 38114, Sept. 20, 1985, as amended at 52 FR 7832, Mar. 13, 1987]

## §520.1010 Furosemide.

(a) *Specifications*. (1) Each tablet contains 12.5 or 50 milligrams (mg) furosemide.

(2) Each bolus contains 2 grams (g) furosemide.

(3) Each packet of powder contains 2 g furosemide.

(4) Each milliliter of syrup contains 10 mg furosemide.

(b) *Sponsors.* See sponsor numbers in §510.600(c) of this chapter for use of dosage forms and strengths listed in paragraph (a) of this section for uses as in paragraph (d) of this section.

(1) No. 000010 for tablets in paragraph (a)(1) of this section for conditions of use in paragraphs (d)(2)(i), (d)(2)(ii)(A), and (d)(3) of this section.

(2) No. 000093 for tablets in paragraph (a)(1) of this section for conditions of use in paragraphs (d)(2)(i) and (d)(2)(ii)(B) of this section.

(3) No. 057926 for tablets in paragraph (a)(1) of this section for conditions of