

PRINCIPLES GOVERNING THE APPLICATION OF
STANDARDS

810.2203 Basis of determination.

GRADES AND GRADE REQUIREMENTS

810.2204 Grades and grade requirements for
wheat.SPECIAL GRADES AND SPECIAL GRADE
REQUIREMENTS810.2205 Special grades and special grade re-
quirements.AUTHORITY: Pub. L. 94-582, 90 Stat. 2867, as
amended (7 U.S.C. 71 *et seq.*).SOURCE: 52 FR 24418, June 30, 1987, unless
otherwise noted.**Subpart A—General Provisions**

NOTE: Compliance with the provisions of these standards does not excuse failure to comply with the provisions of the Federal Food, Drug, and Cosmetic Act, or other Federal laws.

TERMS DEFINED

**§ 810.101 Grains for which standards
are established.**

Grain refers to barley, canola, corn, flaxseed, mixed grain, oats, rye, sorghum, soybeans, sunflower seed, triticale, and wheat. Standards for these food grains, feed grains, and oilseeds are established under the United States Grain Standards Act.

[57 FR 3274, Jan. 29, 1992]

§ 810.102 Definition of other terms.

Unless otherwise stated, the definitions in this section apply to all grains. All other definitions unique to a particular grain are contained in the appropriate subpart for that grain.

(a) *Distinctly low quality.* Grain that is obviously of inferior quality because it is in an unusual state or condition, and that cannot be graded properly by use of other grading factors provided in the standards. Distinctly low quality includes the presence of any objects too large to enter the sampling device; i.e., large stones, wreckage, or similar objects.

(b) *Moisture.* Water content in grain as determined by an approved device according to procedures prescribed in FGIS instructions.

(c) *Stones.* Concreted earthy or mineral matter and other substances of similar hardness that do not disintegrate in water.

(d) *Test weight per bushel.* The weight per Winchester bushel (2,150.42 cubic inches) as determined using an approved device according to procedures prescribed in FGIS instructions. Test weight per bushel in the standards for corn, mixed grain, oats, sorghum, and soybeans is determined on the original sample. Test weight per bushel in the standards for barley, flaxseed, rye, sunflower seed, triticale, and wheat is determined after mechanically cleaning the original sample. Test weight per bushel is recorded to the nearest tenth pound for corn, rye, triticale, and wheat. Test weight per bushel for all other grains, if applicable, is recorded in whole and half pounds with a fraction of a half pound disregarded. Test weight per bushel is not an official factor for canola.

(e) *Whole kernels.* Grain with $\frac{1}{4}$ or less of the kernel removed.

[52 FR 24418, June 30, 1987, as amended at 60
FR 61196, Nov. 29, 1995]PRINCIPLES GOVERNING THE
APPLICATION OF STANDARDS**§ 810.103 Basis of determination.**

(a) *Distinctly low quality.* The determination of distinctly low quality is made on the basis of the lot as a whole at the time of sampling when a condition exists that may or may not appear in the representative sample and/or the sample as a whole.

(b) *Certain quality determinations.* Each determination of rodent pellets, bird droppings, other animal filth, broken glass, castor beans, cockleburrs, crotalaria seeds, dockage, garlic, live insect infestation, large stones, moisture, temperature, an unknown foreign substance(s), and a commonly recognized harmful or toxic substance(s) is made on the basis of the sample as a whole. When a condition exists that may not appear in the representative sample, the determination may be made on the basis of the lot as a whole at the time of sampling according to procedures prescribed in FGIS instructions.