

(2) *Tannin sorghum*. Sorghum which is high in tannin content due to the presence of a pigmented testa (subcoat) and contains not more than 10.0 percent non-Tannin sorghum. The pericarp color of this class is usually brown but may also be white, yellow, pink, orange, red, or bronze.

(3) *White sorghum*. Sorghum which is low in tannin content due to the absence of a pigmented testa (subcoat) and contains not more than 2.0 percent sorghum of other classes. The pericarp color of this class is white or translucent and includes sorghum containing spots that, singly or in combination, cover 25.0 percent or less of the kernel.

(4) *Mixed sorghum*. Sorghum which does not meet the requirements for any of the classes Sorghum, Tannin sorghum, or White sorghum.

(d) *Damaged kernels*. Kernels, pieces of sorghum kernels and other grains that are badly ground damaged, badly weather damaged, diseased, frost-damaged, germ-damaged, heat-damaged, insect-bored, mold-damaged, sprout-damaged, or otherwise materially damaged.

(e) *Dockage*. All matter other than sorghum that can be removed from the original sample by use of an approved device according to procedures prescribed in FGIS instructions. Also, underdeveloped, shriveled, and small pieces of sorghum kernels removed in properly separating the material other than sorghum.

(f) *Foreign material*. All matter, except sorghum, which passes over the number 6 riddle and all matter other than sorghum that remains on top of the 5/64 triangular-hole sieve according to procedures prescribed in FGIS instructions.

(g) *Heat-damaged kernels*. Kernels, pieces of sorghum kernels, and other grains that are materially discolored and damaged by heat.

(h) *Nongrain sorghum*. Seeds of broomcorn, Johnson-grass, *Sorghum almum* Parodi, sorghum-sudangrass hybrids, sorgrass, sudangrass, and sweet sorghum (*sorgo*).

(i) *Pericarp*. The pericarp is the outer layers of the sorghum grain and is fused to the seedcoat.

(j) *Sieves*.

(1) *1.98 mm (5/64 (0.0781) inches) triangular-hole sieve*. A metal sieve 0.81 mm (0.032 inches) thick with equilateral triangular perforations the inscribed circles of which are 1.98 mm (0.0781 inches) in diameter.

(2) *0.99 mm (2 1/2 /64 (0.0391) inches) round-hole sieve*. A metal sieve 0.81 mm (0.032 inch) thick with round holes 0.99 mm (0.0391 inches) in diameter.

[52 FR 24418, June 30, 1987, as amended at 52 FR 24437, June 30, 1987; 52 FR 28534, July 31, 1987; 57 FR 58971, Dec. 14, 1992]

PRINCIPLES GOVERNING THE
APPLICATION OF STANDARDS

§ 810.1403 Basis of determination.

Each determination of broken kernels and foreign material is made on the basis of the grain when free from dockage. Each determination of class, damaged kernels, heat-damaged kernels, and stones is made on the basis of the grain when free from dockage and that portion of the broken kernels, and foreign material that will pass through a 1.98 mm (5/64 inches) triangular-hole sieve. Other determinations not specifically provided for in the general provisions are made on the basis of the grain as a whole except the determination of odor is made on either the basis of the grain as a whole or the grain when free from dockage, broken kernels, and foreign material removed by the 1.98 mm (5/64 inches) triangular-hole sieve.

[57 FR 58971, Dec. 14, 1992]

§ 810.1404

7 CFR Ch. VIII (1–1–02 Edition)

GRADES AND GRADE REQUIREMENTS

§ 810.1404 Grades and grade requirements for sorghum.

Grading factors	Grades U.S. Nos. ¹			
	1	2	3	4
Minimum pound limits of:				
Test weight per bushel	57.0	55.0	53.0	51.0
Maximum percent limits of:				
Damaged kernels:				
Heat (part of total)	0.2	0.5	1.0	3.0
Total	2.0	5.0	10.0	15.0
Broken kernels and foreign material:				
Foreign material (part of total)	1.5	2.5	3.5	4.5
Total	4.0	7.0	10.0	13.0
Maximum count limits of:				
Other material:				
Animal filth	9	9	9	9
Castor beans	1	1	1	1
Crotalaria seeds	2	2	2	2
Glass	1	1	1	1
Stones ²	7	7	7	7
Unknown foreign substance	3	3	3	3
Cockleburrs	7	7	7	7

U.S. Sample grade is Sorghum that:

(a) Does not meet the requirements for U.S. Nos. 1, 2, 3, or 4; or

(b) Has a musty, sour or commercially objectionable foreign odor (except smut odor); or

(c) Is badly weathered, heating or distinctly low quality.

¹ Sorghum which is distinctly discolored shall not grade higher than U.S. No. 3.
² Aggregate weight of stones must also exceed 0.2 percent of sample weight.

[57 FR 58971, Dec. 14, 1992]

SPECIAL GRADES AND SPECIAL GRADE REQUIREMENTS

§ 810.1405 Special grades and special grade requirements.

Smutty sorghum. Sorghum that has kernels covered with smut spores to give a smutty appearance in mass, or that contains 20 or more smut balls in 100 grams of sorghum.

[52 FR 24418, June 30, 1987, as amended at 52 FR 24441, June 30, 1987]

Subpart J—United States Standards for Soybeans

TERMS DEFINED

§ 810.1601 Definition of soybeans.

Grain that consists of 50 percent or more of whole or broken soybeans (*Glycine max* (L.) Merr.) that will not pass through an ³/₆₄ round-hole sieve

and not more than 10.0 percent of other grains for which standards have been established under the United States Grain Standards Act.

§ 810.1602 Definition of other terms.

(a) *Classes.* There are two classes for soybeans: Yellow soybeans and Mixed soybeans.

(1) *Yellow soybeans.* Soybeans that have yellow or green seed coats and which in cross section, are yellow or have a yellow tinge, and may include not more than 10.0 percent of soybeans of other colors.

(2) *Mixed soybeans.* Soybeans that do not meet the requirements of the class Yellow soybeans.

(b) *Damaged kernels.* Soybeans and pieces of soybeans that are badly ground-damaged, badly weather-damaged, diseased, frost-damaged, germ-damaged, heat-damaged, insect-bored, mold-damaged, sprout-damaged,