

skin protectant identified in §352.20(b), the directions for sunscreens in §352.52(d) should be used. For products marketed as a lip protectant or lipstick, §352.52(d)(4) applies.

Subpart D—Testing Procedures

§ 352.70 Standard sunscreen.

(a) *Laboratory validation.* A standard sunscreen shall be used concomitantly in the testing procedures for determining the SPF value of a sunscreen drug product to ensure the uniform evaluation of sunscreen drug products. The standard sunscreen shall be an 8-percent homosalate preparation with a mean SPF value of 4.47 (standard deviation =1.279). In order for the SPF determination of a test product to be considered valid, the SPF of the standard sunscreen must fall within the standard deviation range of the expected SPF (i.e., 4.47 ± 1.279) and the 95-percent confidence interval for the mean SPF must contain the value 4.

(b) *Preparation of the standard homosalate sunscreen.* (1) The standard homosalate sunscreen is prepared from two different preparations (preparation A and preparation B) with the following compositions:

COMPOSITION OF PREPARATION A AND PREPARATION B OF THE STANDARD SUNSCREEN

Ingredients—	Percent by weight
Preparation A	
Lanolin	5.00
Homosalate	8.00
White petrolatum	2.50
Stearic acid	4.00
Propylparaben	0.05
Preparation B	
Methylparaben	0.10
Edetate disodium	0.05
Propylene glycol	5.00
Triethanolamine	1.00
Purified water U.S.P	74.30

(2) Preparation A and preparation B are heated separately to 77 to 82 °C, with constant stirring, until the contents of each part are solubilized. Add preparation A slowly to preparation B while stirring. Continue stirring until the emulsion formed is cooled to room temperature (15 to 30 °C). Add sufficient purified water to obtain 100 grams of standard sunscreen preparation.

(c) *Assay of the standard homosalate sunscreen.* Assay the standard

homosalate sunscreen preparation by the following method to ensure proper concentration:

(1) *Preparation of the assay solvent.* The solvent consists of 1 percent glacial acetic acid (V/V) in denatured ethanol. The denatured ethanol should not contain a UV radiation absorbing denaturant.

(2) *Preparation of a 1-percent solution of the standard homosalate sunscreen preparation.* Accurately weigh 1 gram of the standard homosalate sunscreen preparation into a 100-milliliter volumetric flask. Add 50 milliliters of the assay solvent. Heat on a steam bath and mix well. Cool the solution to room temperature (15 to 30 °C). Then dilute the solution to volume with the assay solvent and mix well to make a 1-percent solution.

(3) *Preparation of the test solution (1:50 dilution of the 1-percent solution).* Filter a portion of the 1-percent solution through number 1 filter paper. Discard the first 10 to 15 milliliters of the filtrate. Collect the next 20 milliliters of the filtrate (second collection). Add 1 milliliter of the second collection of the filtrate to a 50-milliliter volumetric flask. Dilute this solution to volume with assay solvent and mix well. This is the test solution (1:50 dilution of the 1-percent solution).

(4) *Spectrophotometric determination.* The absorbance of the test solution is measured in a suitable double beam spectrophotometer with the assay solvent and reference beam at a wavelength near 306 nanometers.

(5) *Calculation of the concentration of homosalate.* The concentration of homosalate is determined by the following formula which takes into consideration the absorbance of the sample of the test solution, the dilution of the 1-percent solution (1:50), the weight of the sample of the standard homosalate sunscreen preparation (1 gram), and the standard absorbance value (172) of homosalate as determined by averaging the absorbance of a large number of batches of raw homosalate:

$$\text{Concentration of homosalate} = \text{absorbance} \times 50 \times 100 \times 172 = \text{percent concentration by weight.}$$