

produce the first perceptible, unambiguous redness reaction with clearly defined borders at 22 to 24 hours postexposure. The SPF value of the test sunscreen is then calculated from the dose of UV radiation required to produce the MED of the protected skin and from the dose of UV radiation required to produce the MED of the unprotected skin (control site) as follows:

SPF value = the ratio of erythema effective exposure (Joules per square meter) (MED(PS)) to the erythema effective exposure (Joules per square meter) (MED(US)).

(d) *Determination of the test product's SPF value and PCD.* Use data from at least 20 test subjects with n representing the number of subjects used. First, for each subject, compute the SPF value as stated in §352.73(b) and (c). Second, compute the mean SPF value, \bar{x} , and the standard deviation, s , for these subjects. Third, obtain the upper 5-percent point from the t distribution table with $n-1$ degrees of freedom. Denote this value by t . Fourth, compute ts/\sqrt{n} . Denote this quantity by A (i.e., $A = ts/\sqrt{n}$). Fifth, calculate the SPF value to be used in labeling as follows: the label SPF equals the largest whole number less than $\bar{x} - A$. Sixth and last, the drug product is classified into a PCD as follows: if $30 + A < \bar{x}$, the PCD is High; if $12 + A < \bar{x} < 30 + A$, the PCD is Moderate; if $2 + A < \bar{x} < 12 + A$, the PCD is Minimal; if $\bar{x} < 2 + A$, the product shall not be labeled as a sunscreen drug product and shall not display an SPF value.

§352.76 Determination if a product is water resistant or very water resistant.

The general testing procedures in §352.72 shall be used as part of the following tests, except where modified in this section. An indoor fresh water pool, whirlpool, and/or jacuzzi maintained at 23 to 32 °C shall be used in these testing procedures. Fresh water is clean drinking water that meets the standards in 40 CFR part 141. The pool and air temperature and the relative humidity shall be recorded.

(a) *Procedure for testing the water resistance of a sunscreen product.* For sunscreen products making the claim of "water resistant," the label SPF shall

be the label SPF value determined after 40 minutes of water immersion using the following procedure for the water resistance test:

(1) Apply sunscreen product (followed by the waiting period after application of the sunscreen product indicated on the product labeling).

(2) 20 minutes moderate activity in water.

(3) 20-minute rest period (do not towel test sites).

(4) 20 minutes moderate activity in water.

(5) Conclude water test (air dry test sites without toweling).

(6) Begin solar simulator exposure to test site areas as described in §352.73.

(b) *Procedure for testing a very water resistant sunscreen product.* For sunscreen products making the claim of "very water resistant," the label SPF shall be the label SPF value determined after 80 minutes of water immersion using the following procedure for the very water resistant test:

(1) Apply sunscreen product (followed by the waiting period after application of the sunscreen product indicated on the product labeling).

(2) 20 minutes moderate activity in water.

(3) 20-minute rest period (do not towel test sites).

(4) 20 minutes moderate activity in water.

(5) 20-minute rest period (do not towel test sites).

(6) 20 minutes moderate activity in water.

(7) 20-minute rest period (do not towel test sites).

(8) 20 minutes moderate activity in water.

(9) Conclude water test (air dry test sites without toweling).

(10) Begin solar simulator exposure to test site areas as described in §352.73.

§352.77 Test modifications.

The formulation or mode of administration of certain products may require modification of the testing procedures in this subpart. In addition, alternative methods (including automated or in vitro procedures) employing the same basic procedures as those described in this subpart may be used. Any proposed

modification or alternative procedure shall be submitted as a petition in accord with §10.30 of this chapter. The petition should contain data to support the modification or data demonstrating that an alternative procedure provides results of equivalent accuracy. All information submitted will be subject to the disclosure rules in part 20 of this chapter.

PART 355—ANTICARIES DRUG PRODUCTS FOR OVER-THE-COUNTER HUMAN USE

Subpart A—General Provisions

Sec.

355.1 Scope.

355.3 Definitions.

Subpart B—Active Ingredients

355.10 Anticaries active ingredients.

355.20 Packaging conditions.

Subpart C—Labeling

355.50 Labeling of anticaries drug products.

355.55 Principal display panel of all fluoride rinse drug products.

335.60 Professional labeling.

Subpart D—Testing Procedures

355.70 Testing procedures for fluoride dentifrice drug products.

AUTHORITY: 21 U.S.C. 321, 351, 352, 353, 355, 360, 371.

SOURCE: 60 FR 52507, Oct. 6, 1995, unless otherwise noted.

EDITORIAL NOTE: Nomenclature changes to part 355 appear at 69 FR 13717, Mar. 24, 2004.

Subpart A—General Provisions

§355.1 Scope.

(a) An over-the-counter anticaries drug product in a form suitable for topical administration to the teeth is generally recognized as safe and effective and is not misbranded if it meets each condition in this part and each general condition established in §330.1 of this chapter.

(b) References in this part to regulatory sections of the Code of Federal Regulations are to chapter I of title 21 unless otherwise noted.

§355.3 Definitions.

As used in this part:

(a) *Abrasive*. Solid materials that are added to dentifrices to facilitate mechanical removal of dental plaque, debris, and stain from tooth surfaces.

(b) *Anhydrous glycerin*. An ingredient that may be prepared by heating glycerin U.S.P. at 150 °C for 2 hours to drive off the moisture content.

(c) *Anticaries drug*. A drug that aids in the prevention and prophylactic treatment of dental cavities (decay, caries).

(d) *Dental caries*. A disease of calcified tissues of teeth characterized by demineralization of the inorganic portion and destruction of the organic matrix.

(e) *Dentifrice*. An abrasive-containing dosage form (gel, paste, or powder) for delivering an anticaries drug to the teeth.

(f) *Fluoride*. The inorganic form of the chemical element fluorine in combination with other elements.

(g) *Fluoride ion*. The negatively charged atom of the chemical element fluorine.

(h) *Fluoride supplement*. A special treatment rinse dosage form that is intended to be swallowed, and is promoted to health professionals for use in areas where the water supply contains 0 to 0.7 parts per million (ppm) fluoride ion.

(i) *Preventive treatment gel*. A dosage form for delivering an anticaries drug to the teeth. Preventive treatment gels are formulated in an anhydrous glycerin base with suitable thickening agents included to adjust viscosity. Preventive treatment gels do not contain abrasives.

(j) *Treatment rinse*. A liquid dosage form for delivering an anticaries drug to the teeth.

(k) *Treatment rinse concentrated solution*. A fluoride treatment rinse in a concentrated form to be mixed with water before using to result in the appropriate fluoride concentration specified in the monograph.

(l) *Treatment rinse effervescent tablets*. A fluoride treatment rinse prepared by adding an effervescent tablet (a concentrated solid dosage form) to water