

disinfectants or other disinfectant¹ approved specifically for this purpose by the Administrator:

(i) A 5 percent solution of sodium hydroxide or commercial lye containing at least 94 percent of sodium hydroxide. The solution shall be freshly prepared immediately before use by dissolving 2½ pounds of sodium hydroxide or lye in 5½ gallons of hot water and shall be applied as near scalding hot as possible to be most effective. (Owing to the extremely caustic nature of sodium hydroxide solution, precautionary measures such as the wearing of rubber gloves and boots to protect the hands and feet, and goggles to protect the eyes, should be taken by those engaged in the disinfection process. It is also advisable to have an acid solution, such as vinegar, in readiness in case any of the sodium hydroxide solution should come in contact with any part of the body.)

(ii) A solution of sodium hypochlorite containing approximately one-half of 1 percent (5,000 parts per million) of available chlorine. The solution shall be freshly prepared.

(iii) When a disinfectant solution has been applied to equipment which will afterwards contact product, the equipment shall be rinsed with clean water before such contact.

(2) In case anthrax infection is found in the hog slaughtering department, an immediate preliminary disinfection shall be made from the head-dropper's station to the point where the disease is detected and the affected carcasses shall be cut down from the rail and removed from the room. Upon completion of the slaughtering of the lot of hogs of which the anthrax-infected animals were a part, slaughtering operations shall cease, and a thorough cleanup and disinfection shall be made, as provided in paragraph (e)(1) of this section. If the slaughter of the lot has not been completed by the close of the day on which anthrax was detected, the cleanup and disinfection shall not be deferred beyond the close of that day.

¹A list of disinfectants approved for this purpose is available upon request to the Scientific Services, Meat and Poultry Inspection, Food Safety and Inspection Service, U.S. Department of Agriculture, Washington, DC 20250.

(3) The first and indispensable precautionary step for persons who have handled anthrax material is thorough cleansing of the hands and arms with liquid soap and running hot water. It is important that this step be taken immediately after exposure, before vegetative anthrax organisms have had time to form spores. In the cleansing, a brush or other appropriate appliance shall be used to insure the removal of all contaminating material from under and about the fingernails. This process of cleansing is most effective when performed in repeated cycles of lathering and rinsing rather than in spending the same amount of time in scrubbing with a single lathering. After the hands have been cleansed thoroughly and rinsed free of soap, they may, if desired, be immersed for about 1 minute in a 1:1,000 solution of bichloride of mercury, followed by thorough rinsing in clean running water. Supplies of bichloride of mercury for the purpose must be held in the custody of the veterinary medical officer. (As a precautionary measure, all persons exposed to anthrax infection should report promptly any suspicious condition (sore or carbuncle) or symptom to a physician, in order that anti-anthrax serum or other treatment may be administered as indicated.)

[35 FR 15567, Oct. 3, 1970; 36 FR 11903, June 23, 1971]

§310.10 Carcasses with skin or hide on; cleaning before evisceration; removal of larvae of *Hypodermae*, external parasites and other pathological skin conditions.

When a carcass is to be dressed with the skin or hide left on, the skin or hide shall be thoroughly washed and cleaned before any incision is made for the purpose of removing any part thereof or evisceration, except that where calves are slaughtered by the kosher method, the heads shall be removed from the carcasses, before washing of the carcasses. The skin shall be removed at the time of post-mortem inspection from any calf carcass infested with the larvae of the "oxwarble" fly (*Hypoderma lineata* and *Hypoderma bovis*), or external parasites, or affected with other pathological skin conditions.