

### § 311.13

### 9 CFR Ch. III (1-1-06 Edition)

#### § 311.13 Pigmentary conditions; melanosis, xanthosis, ochronosis, etc.

(a) Except as provided in § 311.19, carcasses of livestock showing generalized pigmentary deposits shall be condemned.

(b) The affected parts of carcasses showing localized pigmentary deposits of such character as to be unwholesome or otherwise adulterated shall be removed and condemned.

#### § 311.14 Abrasions, bruises, abscesses, pus, etc.

All slight, well-limited abrasions on the tongue and inner surface of the lips and mouth, when without lymph node involvement, shall be carefully excised, leaving only sound, normal tissue, which may be passed for human food. Any organ or other part of a carcass which is badly bruised or which is affected by an abscess, or a suppurating sore shall be condemned; and when the lesions are of such character or extent as to affect the whole carcass, the whole carcass shall be condemned. Portions of carcasses which are contaminated by pus or other diseased material shall be condemned.

#### § 311.15 Brucellosis.

Carcasses affected with localized lesions of brucellosis may be passed for human food after the affected parts are removed and condemned.

#### § 311.16 Carcasses so infected that consumption of the meat may cause food poisoning.

(a) All carcasses of animals so infected that consumption of the products thereof may give rise to food poisoning shall be condemned. This includes all carcasses showing signs of:

(1) Acute inflammation of the lungs, pleura, pericardium, peritoneum, or meninges.

(2) Septicemia or pyemia, whether puerperal, traumatic, or without any evident cause.

(3) Gangrenous or severe hemorrhagic enteritis or gastritis.

(4) Acute diffuse metritis or mastitis.

(5) Phlebitis of the umbilical veins.

(6) Septic or purulent traumatic pericarditis.

(7) Any acute inflammation, abscess, or suppurating sore, if associated with acute nephritis, fatty and degenerated liver, swollen soft spleen, marked pulmonary hyperemia, general swelling of lymph nodes, diffuse redness of the skin, cachexia, icteric discoloration of the carcass or similar condition, either singly or in combination.

(8) Salmonellosis.

(b) Implements contaminated by contact with carcasses affected with any of the disease conditions mentioned in this section shall be thoroughly cleaned and sanitized as prescribed in part 308 of this subchapter. The equipment used in the dressing of such carcasses, such as viscera trucks or inspection tables, shall be sanitized with hot water having a minimum temperature of 180 °F. Carcasses or parts of carcasses contaminated by contact with such diseased carcasses shall be condemned unless all contaminated tissues are removed within 2 hours.

#### § 311.17 Necrobacillosis, pyemia, and septicemia.

From the standpoint of meat inspection, necrobacillosis may be regarded as a local infection at the beginning, and carcasses in which the lesions are localized may be passed for human food if in a good state of nutrition, after those portions affected with necrotic lesions are removed and condemned. However, when emaciation, cloudy swelling of the parenchymatous tissue of organs or enlargement of the lymph nodes is associated with the infection, it is evident that the disease has progressed beyond the condition of localization to a state of toxemia, and the entire carcass shall therefore be condemned as both unwholesome and noxious. Pyemia or septicemia may intervene as a complication of the local necrosis, and when present the carcass shall be condemned in accordance with § 311.16.

#### § 311.18 Caseous lymphadenitis.

(a) A thin carcass showing well-marked lesions in the viscera and the skeletal lymph nodes, or a thin carcass showing extensive lesions in any part shall be condemned.

(b) A thin carcass showing well-marked lesions in the viscera with only

slight lesions elsewhere or showing well-marked lesions in the skeletal lymph nodes with only slight lesions elsewhere may be passed for cooking.

(c) A thin carcass showing only slight lesions in the skeletal lymph nodes and in the viscera may be passed for human food without restriction.

(d) A well-nourished carcass showing well-marked lesions in the viscera and with only slight lesions elsewhere or showing well-marked lesions confined to the skeletal lymph nodes with only slight lesions elsewhere may be passed for human food without restriction.

(e) A well-nourished carcass showing well-marked lesions in the viscera and the skeletal lymph nodes may be passed for cooking; but where the lesions in a well-nourished carcass are both numerous and extensive, it shall be condemned.

(f) All affected organs and nodes of carcasses passed for human food without restriction or passed for cooking shall be removed and condemned.

(g) As used in this section, the term "thin" does not apply to a carcass which is anemic or emaciated; and the term "lesions" refers to lesions of caseous lymphadenitis.

#### § 311.19 Icterus.

Carcasses showing any degree of icterus shall be condemned. Yellow fat conditions caused by nutritional factors or characteristic of certain breeds of livestock and yellow fat sometimes seen in sheep shall not be confused with icterus. Such carcasses should be passed for human food, if otherwise normal.

#### § 311.20 Sexual odor of swine.

(a) Carcasses of swine which give off a pronounced sexual odor shall be condemned.

(b) The meat of swine carcasses which give off a sexual odor less than pronounced may be passed for use in comminuted cooked meat food product or for rendering. Otherwise it shall be condemned.

#### § 311.21 Mange or scab.

Carcasses of livestock affected with mange or scab in advanced stages, showing cachexia or extensive inflammation of the flesh, shall be con-

demned. When the disease is slight, the carcass may be passed after removal of the affected portion.

#### § 311.22 Hogs affected with urticaria, tinea tonsurans, demodex folliculorum, or erythema.

Carcasses of hogs affected with urticaria (nettle rash), tinea tonsurans, demodex folliculorum, or erythema may be passed for human food after detaching and condemning the affected skin, if the carcass is otherwise not adulterated.

#### § 311.23 Tapeworm cysts (cysticercus bovis) in cattle.

(a) Except as provided in paragraph (b) of this section, carcasses of cattle affected with lesions of cysticercus bovis shall be disposed of as follows:

(1) Carcasses of cattle displaying lesions of cysticercus bovis shall be condemned if the infestation is extensive or if the musculature is edematous or discolored. Carcasses shall be considered extensively infested if in addition to finding lesions in at least two of the usual inspection sites, namely the heart, diaphragm and its pillars, muscles of mastication, esophagus, tongue, and musculature exposed during normal dressing operations, they are found in at least two of the sites exposed by (i) an incision made into each round exposing the musculature in cross section, and (ii) a transverse incision into each forelimb commencing 2 or 3 inches above the point of the olecranon and extending to the humerus.

(2) Carcasses of cattle showing one or more tapeworm lesions of cysticercus bovis but not so extensive as indicated in paragraph (a)(1) of this section, as determined by a careful examination, including examination of, but not limited to, the heart, diaphragm and its pillars, muscles of mastication, esophagus, tongue, and musculature exposed during normal dressing operations, may be passed for human food after removal and condemnation of the lesions with surrounding tissues: *Provided*, That the carcasses, appropriately identified by retained tags, are held in cold storage under positive control of a USDA Food Inspector at a temperature not higher than 15 °F. continuously for a period of not less than 10 days, or in