

§ 421.185

NSPS FOR THE PRIMARY AND SECONDARY GERMANIUM AND GALLIUM SUBCATEGORY

Pollutant or pollutant property	Maximum for any 1 day	Maximum for monthly average
	mg/kg (pounds per million pounds) of germanium hydrolyzed	
Arsenic	39.440	17.550
Lead	7.925	3.774
Zinc	27.550	11.510
Fluoride	660.500	375.500
Total suspended solids	773.700	368.000
pH	(¹)	(¹)

¹ Within the range of 7.5 to 10.0 at all times.

(d) Acid wash and rinse water.

NSPS FOR THE PRIMARY AND SECONDARY GERMANIUM AND GALLIUM SUBCATEGORY

Pollutant or pollutant property	Maximum for any 1 day	Maximum for monthly average
	mg/kg (pounds per million pounds) of germanium washed	
Arsenic	325.50	144.80
Lead	65.40	31.14
Zinc	227.40	94.99
Fluoride	5,450.00	3,099.00
Total suspended solids	6,385.00	3,037.00
pH	(¹)	(¹)

¹ Within the range of 7.5 to 10.0 at all times.

(e) Gallium hydrolysis filtrate.

NSPS FOR THE PRIMARY AND SECONDARY GERMANIUM AND GALLIUM SUBCATEGORY

Pollutant or pollutant property	Maximum for any 1 day	Maximum for monthly average
	mg/kg (pounds per million pounds) of gallium hydrolyzed	
Arsenic	70.450	31.350
Lead	14.160	6.742
Zinc	49.220	20.560
Fluoride	1,180.000	670.800
Total suspended solids	1,382.000	657.300
pH	(¹)	(¹)

¹ Within the range of 7.5 to 10.0 at all times.

(f) Solvent extraction raffinate.

40 CFR Ch. I (7-1-06 Edition)

NSPS FOR THE PRIMARY AND SECONDARY GERMANIUM AND GALLIUM SUBCATEGORY

Pollutant or pollutant property	Maximum for any 1 day	Maximum for monthly average
	mg/kg (pounds per million pounds) of gallium produced by solvent extraction	
Arsenic	39.330	17.500
Lead	7.904	3.764
Zinc	27.480	11.480
Fluoride	658.700	374.500
Total suspended solids	771.600	367.000
pH	(¹)	(¹)

¹ Within the range of 7.5 to 10.0 at all times.

§ 421.185 Pretreatment standards for existing sources.

Except as provided in 40 CFR 403.7 and 403.13, any existing source subject to this subpart which introduces pollutants into a publicly owned treatment works must comply with 40 CFR part 403 and achieve the following pretreatment standards for existing sources. The mass of wastewater pollutants in primary and secondary germanium and gallium process wastewater introduced into a POTW must not exceed the following values:

(a) Still liquor.

PSES FOR THE PRIMARY AND SECONDARY GERMANIUM AND GALLIUM SUBCATEGORY

Pollutant or pollutant property	Maximum for any 1 day	Maximum for monthly average
	mg/kg (pounds per million pounds) of germanium chlorinated	
Arsenic	131.70	58.59
Lead	26.46	12.60
Zinc	91.98	38.43
Fluoride	2,205.00	1,254.00

(b) Chlorinator wet air pollution control.

PSES FOR THE PRIMARY AND SECONDARY GERMANIUM AND GALLIUM SUBCATEGORY

Pollutant or pollutant property	Maximum for any 1 day	Maximum for monthly average
	mg/kg (pounds per million pounds) of germanium chlorinated	
Arsenic	27.530	12.250
Lead	5.531	2.634
Zinc	19.230	8.034

Environmental Protection Agency

§ 421.186

PSES FOR THE PRIMARY AND SECONDARY GERMANIUM AND GALLIUM SUBCATEGORY—Continued

Pollutant or pollutant property	Maximum for any 1 day	Maximum for monthly average
Fluoride	461.000	262.100

(c) Germanium hydrolysis filtrate.

PSES FOR THE PRIMARY AND SECONDARY GERMANIUM AND GALLIUM SUBCATEGORY

Pollutant or pollutant property	Maximum for any 1 day	Maximum for monthly average
	mg/kg (pounds per million pounds) of germanium hydrolyzed	
Arsenic	39.440	17.550
Lead	7.925	3.774
Zinc	27.550	11.510
Fluoride	660.500	375.500

(d) Acid wash and rinse water.

PSES FOR THE PRIMARY AND SECONDARY GERMANIUM AND GALLIUM SUBCATEGORY

Pollutant or pollutant property	Maximum for any 1 day	Maximum for monthly average
	mg/kg (pounds per million pounds) of germanium washed	
Arsenic	325.50	144.80
Lead	65.40	31.14
Zinc	227.40	94.99
Fluoride	5,450.00	3,099.00

(e) Gallium hydrolysis filtrate.

PSES FOR THE PRIMARY AND SECONDARY GERMANIUM AND GALLIUM SUBCATEGORY

Pollutant or pollutant property	Maximum for any 1 day	Maximum for monthly average
	mg/kg (pounds per million pounds) of gallium hydrolyzed	
Arsenic	70.450	31.350
Lead	14.160	6.742
Zinc	49.220	20.560
Fluoride	1,180.000	670.800

(f) Solvent extraction raffinate.

PSES FOR THE PRIMARY AND SECONDARY GERMANIUM AND GALLIUM SUBCATEGORY

Pollutant or pollutant property	Maximum for any 1 day	Maximum for monthly average
	mg/kg (pounds per million pounds) of gallium produced by solvent extraction	
Arsenic	39.330	17.500
Lead	7.904	3.764
Zinc	27.480	11.480
Fluoride	658.700	374.500

§ 421.186 Pretreatment standards for new sources.

Except as provided in 40 CFR 403.7, any new source subject to this subpart which introduces pollutants into a publicly owned treatment works must comply with 40 CFR part 403 and achieve the following pretreatment standards for new sources. The mass of wastewater pollutants in primary and secondary germanium and gallium process wastewater introduced into a POTW shall not exceed the following values:

(a) Still Liquor.

PSNS FOR THE PRIMARY AND SECONDARY GERMANIUM AND GALLIUM SUBCATEGORY

Pollutant or pollutant property	Maximum for any 1 day	Maximum for monthly average
	mg/kg (pounds per million pounds) of germanium chlorinated	
Arsenic	131.70	58.59
Lead	26.46	12.60
Zinc	91.98	38.43
Fluoride	2,205.00	1,254.00

(b) Chlorinator Wet Air Pollution Control.

PSNS FOR THE PRIMARY AND SECONDARY GERMANIUM AND GALLIUM SUBCATEGORY

Pollutant or pollutant property	Maximum for any 1 day	Maximum for monthly average
	mg/kg (pounds per million pounds) of germanium chlorinated	
Arsenic	27.530	12.250
Lead	5.531	2.634
Zinc	19.230	8.034
Fluoride	461.000	262.100

(c) Germanium Hydrolysis Filtrate.