

terms of the minimum dosage recommended per minimum time interval.

(2) May contain an indication for the symptomatic relief of hyperacidity associated with the diagnosis of peptic ulcer, gastritis, peptic esophagitis, gastric hyperacidity, and hiatal hernia.

(3) *For products containing basic aluminum carbonate gel identified in § 331.11(a)(1)—Indication.* “For the treatment, control, or management of hyperphosphatemia, or for use with a low phosphate diet to prevent formation of phosphate urinary stones, through the reduction of phosphates in the serum and urine.”

(4) *For products containing aluminum identified in § 331.11(a)—Warnings.* (i) Prolonged use of aluminum-containing antacids in patients with renal failure may result in or worsen dialysis osteomalacia. Elevated tissue aluminum levels contribute to the development of the dialysis encephalopathy and osteomalacia syndromes. Small amounts of aluminum are absorbed from the gastrointestinal tract and renal excretion of aluminum is impaired in renal failure. Aluminum is not well removed by dialysis because it is bound to albumin and transferrin, which do not cross dialysis membranes. As a result, aluminum is deposited in bone, and dialysis osteomalacia may develop when large amounts of aluminum are ingested orally by patients with impaired renal function.

(ii) Aluminum forms insoluble complexes with phosphate in the gastrointestinal tract, thus decreasing phosphate absorption. Prolonged use of aluminum-containing antacids by normophosphatemic patients may result in hypophosphatemia if phosphate intake is not adequate. In its more severe forms, hypophosphatemia can lead to anorexia, malaise, muscle weakness, and osteomalacia.

(b) Professional labeling for an antacid-antiflatulent combination may contain the information allowed for health professionals for antacids and antiflatulents.

[39 FR 19874, June 4, 1974. Redesignated and amended at 55 FR 19859, May 11, 1990]

PART 332—ANTIFLATULENT PRODUCTS FOR OVER-THE-COUNTER HUMAN USE

Subpart A—General Provisions

Sec.

332.1 Scope.

332.3 Definitions.

Subpart B—Active Ingredients

332.10 Antiflatulent active ingredients.

332.15 Combination with non-antiflatulent active ingredients.

Subpart C—Labeling

332.30 Labeling of antiflatulent products.

332.31 Professional labeling.

AUTHORITY: 21 U.S.C. 321, 351, 352, 353, 355, 360, 371.

SOURCE: 39 FR 19877, June 4, 1974, unless otherwise noted.

Subpart A—General Provisions

§ 332.1 Scope.

An over-the-counter antiflatulent product in a form suitable for oral administration is generally recognized as safe and effective and is not misbranded if it meets each of the following conditions and each of the general conditions established in § 330.1 of this chapter.

§ 332.3 Definitions.

As used in this part:

Antigas. A term that may be used interchangeably with the term antiflatulent. Neither term should be considered as describing the mechanism of action of the active ingredient contained in the product.

[61 FR 8838, Mar. 5, 1996]

Subpart B—Active Ingredients

§ 332.10 Antiflatulent active ingredients.

Simethicone; maximum daily dose 500 mg. There is no dosage limitation at this time for professional labeling.