

Blood or Red Blood Cells intended for autologous infusion:

(1) Information adequately identifying the patient, e.g., name, blood group, hospital, and identification number.

(2) Date of donation.

(3) The statement: "FOR AUTOLOGOUS USE ONLY."

(4) In place of the blood group label, each container of blood intended for autologous use and obtained from a donor who fails to meet any of the donor suitability requirements under § 640.3 of this chapter or who is reactive in the hepatitis tests prescribed under § 610.40 of this chapter shall be prominently and permanently labeled: "FOR AUTOLOGOUS USE ONLY."

(5) Units of blood originally intended for autologous use, except those labeled as prescribed under paragraph (i)(4) of this section, may be issued for homologous transfusion provided the container label complies with all applicable provisions of paragraphs (b) through (e) of this section. In such case, the special label required under paragraph (i) (1), (2), and (3) of this section shall be removed or otherwise obscured.

(j) A tie-tag attached to the container may be used for providing the information required by paragraph (e) (1)(iii), (2)(ii), and (4), (h), or (i)(1), (2), and (3) of this section.

[50 FR 35469, Aug. 30, 1985, as amended at 53 FR 116, Jan. 5, 1988; 55 FR 11014, Mar. 26, 1990; 57 FR 10814, Mar. 31, 1992; 59 FR 23636, May 6, 1994; 63 FR 16685, Apr. 6, 1998; 64 FR 45371, Aug. 19, 1999; 66 FR 31162, June 11, 2001; 67 FR 4907, Feb. 1, 2002; 69 FR 9171, Feb. 26, 2004; 70 FR 14984, Mar. 24, 2005]

§ 606.122 Instruction circular.

An instruction circular shall be available for distribution if the product is intended for transfusion. The instruction circular shall provide adequate directions for use, including the following information:

(a) Instructions to mix the product before use.

(b) Instructions to use a filter in the administration equipment.

(c) The statement "Do Not Add Medications" or an explanation concerning allowable additives.

(d) A description of the product, its source, and preparation, including the name and proportion of the anticoagulant used in collecting the Whole Blood from each product is prepared.

(e) Statements that the product was prepared from blood that was negative when tested for antibody to Human Immunodeficiency Virus (HIV) and non-reactive for hepatitis B surface antigen by FDA required tests and nonreactive when tested for syphilis by a serologic test for syphilis (STS).

(f) The statements: "Warning. The risk of transmitting infectious agents is present. Careful donor selection and available laboratory tests do not eliminate the hazard."

(g) The names of cryoprotective agents and other additives that may still be present in the product.

(h) The names and results of all tests performed when necessary for safe and effective use.

(i) The use of the product, indications, contraindications, side effects and hazards, dosage and administration recommendations.

(j) [Reserved]

(k) For Red Blood Cells, the instruction circular shall contain:

(1) Instructions to administer a suitable plasma volume expander if Red Blood Cells are substituted when Whole Blood is the indicated product.

(2) A warning not to add Lactated Ringer's Injection U.S.P. solution to Red Blood Cell products.

(1) For Platelets, the instruction circular shall contain:

(1) The approximate volume of plasma from which a sample unit of Platelets is prepared.

(2) Instructions to begin administration as soon as possible, but not more than 4 hours after entering the container.

(m) For Plasma, the instruction circular shall contain:

(1) A warning against further processing of the frozen product if there is evidence of breakage or thawing.

(2) Instructions to thaw the frozen product at a temperature between 30 and 37 °C.

(3) When applicable, instructions to begin administration of the product within 6 hours after thawing.

§ 606.140

21 CFR Ch. I (4–1–07 Edition)

(4) Instructions to administer to ABO-group-compatible recipients.

(5) A statement that this product has the same hepatitis risk as Whole Blood; other plasma volume expanders without this risk are available for treating hypovolemia.

(n) For Cryoprecipitated AHF, the instruction circular shall contain:

(1) A statement that the average potency is 80 or more International Units of antihemophilic factor.

(2) The statement: “Usually contains at least 150 milligrams of fibrinogen”; or, alternatively, the average fibrinogen level determined by assay of representative units.

(3) A warning against further processing of the product if there is evidence of breakage or thawing.

(4) Instructions to thaw the product for no more than 15 minutes at a temperature of between 30 and 37 °C.

(5) Instructions to store at room temperature after thawing and to begin administration as soon as possible but no more than 4 hours after entering the container or after pooling and within 6 hours after thawing.

(6) A statement that 0.9 percent Sodium Chloride Injection U.S.P. is the preferred diluent.

(7) Adequate instructions for pooling to ensure complete removal of all concentrated material from each container.

(8) The statement: “Good patient management requires monitoring treatment responses to Cryoprecipitated AHF transfusions with periodic plasma factor VIII or fibrinogen assays in hemophilia A and hypofibrinogenemic recipients, respectively.”

[50 FR 35470, Aug. 30, 1985, as amended at 53 FR 116, Jan. 5, 1988; 64 FR 45371, Aug. 19, 1999]

Subpart H—Laboratory Controls

§ 606.140 Laboratory controls.

Laboratory control procedures shall include:

(a) The establishment of scientifically sound and appropriate specifications, standards and test procedures to assure that blood and blood components are safe, pure, potent and effective.

(b) Adequate provisions for monitoring the reliability, accuracy, precision and performance of laboratory test procedures and instruments.

(c) Adequate identification and handling of all test samples so that they are accurately related to the specific unit of product being tested, or to its donor, or to the specific recipient, where applicable.

§ 606.151 Compatibility testing.

Standard operating procedures for compatibility testing shall include the following:

(a) A method of collecting and identifying the blood samples of recipients to ensure positive identification.

(b) The use of fresh recipient serum or plasma samples less than 3 days old for all pretransfusion testing if the recipient has been pregnant or transfused within the previous 3 months.

(c) Procedures to demonstrate incompatibility between the donor’s cell type and the recipient’s serum or plasma type.

(d) A provision that, if the unit of donor’s blood has not been screened by a method that will demonstrate agglutinating, coating and hemolytic antibodies, the recipient’s cells shall be tested with the donor’s serum (minor crossmatch) by a method that will so demonstrate.

(e) Procedures to expedite transfusion in life-threatening emergencies. Records of all such incidents shall be maintained, including complete documentation justifying the emergency action, which shall be signed by a physician.

[40 FR 53532, Nov. 18, 1975, as amended at 64 FR 45371, Aug. 19, 1999; 66 FR 1835, Jan. 10, 2001; 66 FR 40889, Aug. 6, 2001]

Subpart I—Records and Reports

§ 606.160 Records.

(a)(1) Records shall be maintained concurrently with the performance of each significant step in the collection, processing, compatibility testing, storage and distribution of each unit of blood and blood components so that all steps can be clearly traced. All records shall be legible and indelible, and shall identify the person performing the