

§ 180.123a

40 CFR Ch. I (7-1-07 Edition)

concentration of residues of inorganic bromides from fumigation of the grains with methyl bromide.

(b) *Section 18 emergency exemptions.* [Reserved]

(c) *Tolerances with regional registrations.* A tolerance with regional registration, as defined in §180.1(n), is established for residues of inorganic bromides (calculated as Br) in or on the following food commodity grown in soil fumigated with methyl bromide.

Commodity	Parts per million
Ginger, roots, postharvest	100

(d) *Indirect or inadvertent residues.* [Reserved]

[71 FR 74812, Dec. 13, 2006]

§ 180.123a Inorganic bromide residues in peanut hay and peanut hulls; statement of policy.

(a) Investigations by the Food and Drug Administration show that peanut hay and peanut shells have been used as feed for meat and dairy animals. While many growers now harvest peanuts with combines and leave the hay on the ground to be incorporated into the soil, some growers follow the practice of curing peanuts on the vines in a stack and save the hay for animal feed. Peanut shells or hulls have been used to a minor extent as roughage for cattle feed. It has been established that the feeding to cattle of peanut hay and peanut hulls containing residues of inorganic bromides will contribute considerable residues of inorganic bromides to the meat and milk.

(b) There are no tolerances for inorganic bromides in meat and milk to cover residues from use of such peanut hulls as animal feed. Peanut hulls containing residues of inorganic bromides from the use of methyl bromide are unsuitable as an ingredient in the feed of meat and dairy animals and should not be represented, sold, or used for that purpose.

[58 FR 65555, Dec. 15, 1993]

§ 180.127 Piperonyl butoxide; tolerances for residues.

(a) *General.* (1) Tolerances for residues of the insecticide piperonyl butoxide [(butyl carbityl)(6-propyl

piperonyl)ether] are established in or on the following food commodities:

Commodity	Parts per million
Almond, postharvest	8
Apple, postharvest	8
Barley, postharvest	20
Bean, postharvest	8
Birdseed, mixtures, postharvest	20
Blackberry, postharvest	8
Blueberry, postharvest	8
Boysenberry, postharvest	8
Buckwheat, grain, postharvest	20
Cattle, fat	0.1
Cattle, meat byproducts	0.1
Cattle, meat	0.1
Cherry, sweet, postharvest	8
Cherry, tart, postharvest	8
Cocoa bean, roasted bean, postharvest	8
Coconut, copra, postharvest	8
Corn, field, grain, postharvest	20
Corn, pop, postharvest	20
Cotton, undelinted seed, postharvest	8
Crabapple, postharvest	8
Currant, postharvest	8
Dewberry, postharvest	8
Egg	1
Fig, postharvest	8
Flax, seed, postharvest	8
Goat, fat	0.1
Goat, meat byproducts	0.1
Goat, meat	0.1
Gooseberry, postharvest	8
Grape, postharvest	8
Guava, postharvest	8
Hog, fat	0.1
Hog, meat byproducts	0.1
Hog, meat	0.1
Horse, fat	0.1
Horse, meat byproducts	0.1
Horse, meat	0.1
Loganberry, postharvest	8
Mango, postharvest	8
Milk, fat	0.25
Muskmelon, postharvest	8
Oat, postharvest	8
Orange, postharvest	8
Peach, postharvest	8
Peanut, postharvest	8
Pea, postharvest	8
Pear, postharvest	8
Pineapple, postharvest	8
Plum, prune, fresh, postharvest	8
Potato, postharvest	0.25
Poultry, fat	3
Poultry, meat byproducts	3
Poultry, meat	3
Raspberry, postharvest	8
Rice, postharvest	20
Rye, postharvest	20
Sheep, fat	0.1
Sheep, meat byproducts	0.1
Sheep, meat	0.1
Sorghum, grain, postharvest	8
Sweet potato, postharvest	0.25
Tomato, postharvest	8
Walnut, postharvest	8
Wheat, postharvest	20

(2) Piperonyl butoxide may be safely used in accordance with the following prescribed conditions:

Environmental Protection Agency

§ 180.127

(i) It is used or intended for use in combination with pyrethrins for control of insects:

(A) In cereal grain mills and in storage areas for milled cereal grain products, whereby the amount of piperonyl butoxide is at least equal to but not more than 10 times the amount of pyrethrins in the formulation.

(B) On the outer ply of multiwall paper bags of 50 pounds or more capacity in amounts not exceeding 60 milligrams per square foot, whereby the amount of piperonyl butoxide is equal to 10 times the amount of pyrethrins in the formulation. Such treated bags are to be used only for food, dried.

(C) On cotton bags of 50 pounds or more capacity in amounts not exceeding 55 milligrams per square foot of cloth, whereby the amount of piperonyl butoxide is equal to 10 times the amount of pyrethrins in the formulation. Such treated bags are constructed with waxed paper liners and are to be used only for food, dried that contain 4 percent fat or less.

(D) In two-ply bags consisting of cellophane/polyolefin sheets bound together by an adhesive layer when it is incorporated in the adhesive. The treated sheets shall contain not more than 50 milligrams of piperonyl butoxide per square foot (538 milligrams per square meter). Such treated bags are to be used only for packaging plum, prune, dried; grape, raisin; and other fruit, dried and are to have a maximum ratio of 3.12 milligrams of piperonyl butoxide per ounce of fruit (0.10 milligram of piperonyl butoxide per gram of product).

(E) In food processing and food storage areas: Provided, That the food is removed or covered prior to such use.

(ii) It is used or intended for use in combination with pyrethrins and N-octylbicycloheptene dicarboximide for insect control in accordance with 21 CFR 178.3730.

(iii) A tolerance of 10 parts per million is established for residues of piperonyl butoxide in or on:

(A) Grain, cereal, milled fractions when present therein as a result of its use in cereal grain mills and in storage areas for milled cereal grain products.

(B) Food, dried when present as a result of migration from its use on the

outer ply of multiwall paper bags of 50 pounds or more capacity.

(C) Food treated in accordance with 21 CFR 178.3730.

(D) Food, dried that contain 4 percent fat, or less, when present as a result of migration from its use on the cloth of cotton bags of 50 pounds or more capacity constructed with waxed paper liners.

(E) Food treated in accordance with paragraph (a)(2)(i)(D) and (E) of this section.

(iv) To assure safe use of the pesticide, its label and labeling shall conform to that registered with the U.S. Environmental Protection Agency, and it shall be used in accordance with such label and labeling.

(v) Where tolerances are established on both raw agricultural commodities and processed food made therefrom, the total residues of piperonyl butoxide in or on the processed food shall not be greater than that permitted by the larger of the two tolerances.

(3) Piperonyl butoxide may be safely used in accordance with the following prescribed conditions:

(i) It is used or intended for use in combination with pyrethrins for control of insects:

(A) On the outer ply of multiwall paper bags of 50 pounds or more capacity in amounts not exceeding 60 milligrams per square foot.

(B) On cotton bags of 50 pounds or more capacity in amounts not exceeding 55 milligrams per square foot of cloth. Such treated bags are constructed with waxed paper liners and are to be used only for feed, dried that contain 4 percent fat or less.

(ii) It is used in combination with pyrethrins, whereby the amount of piperonyl butoxide is equal to 10 times the amount of pyrethrins in the formulation. Such treated bags are to be used only for feed, dried.

(iii) A tolerance of 10 parts per million is established for residues of piperonyl butoxide when present as the result of migration:

(A) In or on feed, dried from its use on the outer ply of multiwall paper bags of 50 pounds or more capacity.

(B) In or on feed, dried that contain 4 percent fat, or less, from its use on

§ 180.128

40 CFR Ch. I (7-1-07 Edition)

cotton bags of 50 pounds or more capacity constructed with waxed paper liners.

(iv) To assure safe use of the pesticide, its label and labeling shall conform to that registered with the U.S. Environmental Protection Agency.

(v) Where tolerances are established on both the raw agricultural commodities and food, processed made therefrom, the total residues of piperonyl butoxide in or on the processed food shall not be greater than that permitted by the larger of the two tolerances.

(b) *Section 18 emergency exemptions.* [Reserved]

(c) *Tolerances with regional registrations.* [Reserved]

(d) *Indirect or inadvertent residues.* [Reserved]

[71 FR 74813, Dec. 13, 2006]

§ 180.128 Pyrethrins; tolerances for residues.

(a) *General.* (1) Tolerances for residues of the insecticide pyrethrins (insecticidally active principles of *Chrysanthemum cinerariaefolium*) are established in or on the following food commodities:

Commodity	Parts per million
Almond, postharvest	1
Apple, postharvest	1
Barley, postharvest	3
Bean, postharvest	1
Birdseed, mixtures, postharvest	3
Blackberry, postharvest	1
Blueberry, postharvest	1
Boysenberry, postharvest	1
Buckwheat, grain, postharvest	3
Cattle, fat	0.1(N)
Cattle, meat byproducts	0.1(N)
Cattle, meat	0.1(N)
Cherry, sweet, postharvest	1
Cherry, tart, postharvest	1
Cacao bean, roasted bean, postharvest	1
Coconut, copra, postharvest	1
Corn, field, grain, postharvest	3
Corn, pop, grain, postharvest	3
Cotton, undelinted seed, postharvest	1
Crabapple, postharvest	1
Currant, postharvest	1
Dewberry, postharvest	1
Egg	0.1(N)
Fig, postharvest	1
Flax, seed, postharvest	1
Goat, fat	0.1(N)
Goat, meat byproducts	0.1(N)
Goat, meat	0.1(N)
Gooseberry, postharvest	1
Grape, postharvest	1
Guava, postharvest	1
Hog, fat	0.1(N)

Commodity	Parts per million
Hog, meat byproducts	0.1(N)
Hog, meat	0.1(N)
Horse, fat	0.1(N)
Horse, meat byproducts	0.1(N)
Horse, meat	0.1(N)
Loganberry, postharvest	1
Mango, postharvest	1
Milk, fat	0.5
Muskmelon, postharvest	1
Oat, grain, postharvest	1
Orange, postharvest	1
Peach, postharvest	1
Peanut, postharvest	1
Pea, postharvest	1
Pear, postharvest	1
Pineapple, postharvest	1
Plum, prune, fresh, postharvest	1
Potato, postharvest	0.05
Poultry, fat	0.2
Poultry, meat byproducts	0.2
Poultry, meat	0.2
Raspberry, postharvest	1
Rice, grain, postharvest	3
Rye, postharvest	3
Sheep, fat	0.1(N)
Sheep, meat byproducts	0.1(N)
Sheep, meat	0.1(N)
Sorghum, grain, grain, postharvest	1
Sweet potato, postharvest	0.05
Tomato, postharvest	1
Walnut, postharvest	1
Wheat, postharvest	3

(2) Pyrethrins may be safely used in accordance with the following prescribed conditions:

(i) It is used or intended for use in combination with piperonyl butoxide for control of insects:

(A) In cereal grain mills and in storage areas for milled cereal grain products, whereby the amount of pyrethrins is from 10 percent to 100 percent of the amount of piperonyl butoxide in the formulation.

(B) On the outer ply of multiwall paper bags of 50 pounds or more capacity in amounts not exceeding 6 milligrams per square foot, whereby the amount of pyrethrins is equal to 10 percent of the amount of piperonyl butoxide in the formulation. Such treated bags are to be used only for food, dried.

(C) On cotton bags of 50 pounds or more capacity in amounts not exceeding 5.5 milligrams per square foot of cloth, whereby the amount of pyrethrins is equal to 10 percent of the amount of piperonyl butoxide in the formulation. Such treated bags are constructed with waxed paper liners and are to be used only for food, dried that contain 4 percent fat or less.