

### § 310.515

### 21 CFR Ch. I (4–1–08 Edition)

(2) A statement that, if additive drugs are introduced into the parenteral system, aseptic techniques should be used and the solution should be thoroughly mixed.

(3) A statement that a solution containing an additive drug should not be stored.

(d) This section does not apply to a biological product licensed under the Public Health Service Act of July 1, 1944 (42 U.S.C. 201).

[62 FR 12084, Mar. 14, 1997]

#### § 310.515 Patient package inserts for estrogens.

(a) *Requirement for a patient package insert.* FDA concludes that the safe and effective use of drug products containing estrogens requires that patients be fully informed of the benefits and risks involved in the use of these drugs. Accordingly, except as provided in paragraph (e) of this section, each estrogen drug product restricted to prescription distribution, including products containing estrogens in fixed combinations with other drugs, shall be dispensed to patients with a patient package insert containing information concerning the drug's benefits and risks. An estrogen drug product that does not comply with the requirements of this section is misbranded under section 502(a) of the Federal Food, Drug, and Cosmetic Act.

(b) *Distribution requirements.* (1) For estrogen drug products, the manufacturer and distributor shall provide a patient package insert in or with each package of the drug product that the manufacturer or distributor intends to be dispensed to a patient.

(2) In the case of estrogen drug products in bulk packages intended for multiple dispensing, and in the case of injectables in multiple-dose vials, a sufficient number of patient labeling pieces shall be included in or with each package to assure that one piece can be included with each package or dose dispensed or administered to every patient. Each bulk package shall be labeled with instructions to the dispenser to include one patient labeling piece with each package dispensed or, in the case of injectables, with each dose administered to the patient. This section does not preclude the manufac-

turer or labeler from distributing additional patient labeling pieces to the dispenser.

(3) Patient package inserts for estrogens dispensed in acute-care hospitals or long-term care facilities will be considered to have been provided in accordance with this section if provided to the patient before administration of the first estrogen and every 30 days thereafter, as long as the therapy continues.

(c) *Patient package insert contents.* A patient package insert for an estrogen drug product is required to contain the following information:

- (1) The name of the drug.
- (2) The name and place of business of the manufacturer, packer, or distributor.
- (3) A statement regarding the benefits and proper uses of estrogens.
- (4) The contraindications to use, i.e., when estrogens should not be used.
- (5) A description of the most serious risks associated with the use of estrogens.
- (6) A brief summary of other side effects of estrogens.
- (7) Instructions on how a patient may reduce the risks of estrogen use.
- (8) The date, identified as such, of the most recent revision of the patient package insert.

(d) *Guidance language.* The Food and Drug Administration issues informal labeling guidance texts under § 10.90(b)(9) of this chapter to provide assistance in meeting the requirements of paragraph (c) of this section. Requests for a copy of the guidance text should be directed to the Center for Drug Evaluation and Research, Division of Metabolism and Endocrine Drug Products (HFD-510), Food and Drug Administration, 5600 Fishers Lane, Rockville, MD 20857.

(e) *Exemptions.* This section does not apply to estrogen-progestogen oral contraceptives. Labeling requirements for these products are set forth in § 310.501.

(f) *Requirement to supplement approved application.* Holders of approved applications for estrogen drug products that are subject to the requirements of this section must submit supplements under § 314.70(c) of this chapter to provide for the labeling required by paragraph (a) of this section. Such labeling

may be put into use without advance approval by the Food and Drug Administration.

[55 FR 18723, May 4, 1990]

**§ 310.517 Labeling for oral hypoglycemic drugs of the sulfonylurea class.**

(a) The University Group Diabetes Program clinical trial has reported an association between the administration of tolbutamide and increased cardiovascular mortality. The Food and Drug Administration has concluded that this reported association provides adequate basis for a warning in the labeling. In view of the similarities in chemical structure and mode of action, the Food and Drug Administration also believes it is prudent from a safety standpoint to consider that the possible increased risk of cardiovascular mortality from tolbutamide applies to all other sulfonylurea drugs as well. Therefore, the labeling for oral hypoglycemic drugs of the sulfonylurea class shall include a warning concerning the possible increased risk of cardiovascular mortality associated with such use, as set forth in paragraph (b) of this section.

(b) Labeling for oral hypoglycemic drugs of the sulfonylurea class shall include in boldface type at the beginning of the "Warnings" section of the labeling the following statement:

**SPECIAL WARNING ON INCREASED RISK OF CARDIOVASCULAR MORTALITY**

The administration of oral hypoglycemic drugs has been reported to be associated with increased cardiovascular mortality as compared to treatment with diet alone or diet plus insulin. This warning is based on the study conducted by the University Group Diabetes Program (UGDP), a long-term prospective clinical trial designed to evaluate the effectiveness of glucose-lowering drugs in preventing or delaying vascular complications in patients with non-insulin-dependent diabetes. The study involved 823 patients who were randomly assigned to one of four treatment groups (*Diabetes*, 19 (supp. 2): 747-830, 1970).

UGDP reported that patients treated for 5 to 8 years with diet plus a fixed dose of tolbutamide (1.5 grams per day) had a rate of cardiovascular mortality approximately 2½ times that of patients treated with diet alone. A significant increase in total mortality was not observed, but the use of tolbutamide was discontinued based on the in-

crease in cardiovascular mortality, thus limiting the opportunity for the study to show an increase in overall mortality. Despite controversy regarding the interpretation of these results, the findings of the UGDP study provide an adequate basis for this warning. The patient should be informed of the potential risks and advantages of (name of drug) and of alternative modes of therapy.

Although only one drug in the sulfonylurea class (tolbutamide) was included in this study, it is prudent from a safety standpoint to consider that this warning may also apply to other oral hypoglycemic drugs in this class, in view of their close similarities in mode of action and chemical structure.

[49 FR 14331, Apr. 11, 1984]

**§ 310.518 Drug products containing iron or iron salts.**

Drug products containing elemental iron or iron salts as an active ingredient in solid oral dosage form, e.g., tablets or capsules shall meet the following requirements:

(a) *Labeling.* (1) The label of any drug in solid oral dosage form (e.g., tablets or capsules) that contains iron or iron salts for use as an iron source shall bear the following statement:

WARNING: Accidental overdose or iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctor or poison control center immediately.

(2)(i) The warning statement required by paragraph (a)(1) of this section shall appear prominently and conspicuously on the information panel of the immediate container label.

(ii) If a drug product is packaged in unit-dose packaging, and if the immediate container bears labeling but not a label, the warning statement required by paragraph (a)(1) of this section shall appear prominently and conspicuously on the immediate container labeling in a way that maximizes the likelihood that the warning is intact until all of the dosage units to which it applies are used.

(3) Where the immediate container is not the retail package, the warning statement required by paragraph (a)(1) of this section shall also appear prominently and conspicuously on the information panel of the retail package label.