

Food and Drug Administration, HHS

§ 369.20

§ 310.201(a) of this chapter, or the provisions of an effective application pursuant to section 505(b) of the act.

§ 369.4 Warnings suggested for drugs by formal or informal statements of policy.

The warning and caution statements included in subpart B of this part in no way affect any warning statement suggested for such drugs or devices by any statement of policy or interpretation in subchapter C of this chapter.

[39 FR 11745, Mar. 29, 1974, as amended at 40 FR 13496, Mar. 27, 1975]

§ 369.6 [Reserved]

§ 369.7 Warnings required by official compendia.

Any drug included in the official compendia defined by the act shall bear such warning or caution statement as may be required by such compendia, and no statement in subpart B or subpart C of this part is intended to alter, modify, or permit the omission of any such statement required by such compendia.

§ 369.8 Warning statements in relation to conditions for use.

The mention in any warning or caution statement included in subparts A, B, and C of this part, of a disease condition does not imply a finding on the part of the Food and Drug Administration that any drug or device is efficacious in such condition; nor is any drug or device bearing labeling referring to such disease condition precluded from regulatory action under the applicable provisions of the act if such claim is considered to be misbranding.

§ 369.9 General warnings re accidental ingestion by children.

Section 369.20 includes under certain items, but not all medicines, the statement: "Keep this and all medicines out of children's reach. In case of overdose, get medical help or contact a Poison Control Center right away," or "Keep out of reach of children." However, in view of the possibility of accidental ingestion of drugs, it is not only suggested but is recommended that one of

these statements be used on the label of all drug products.

[64 FR 13296, Mar. 17, 1999]

§ 369.10 Conspicuousness of warning statements.

Necessary warning statements should appear in the labeling prominently and conspicuously as compared to other words, statements, designs, and devices, and in bold type on clearly contrasting background, in order to comply with the provisions of section 502(c) and (f)(2) of the act. The warning statements should be placed in the labeling in juxtaposition with the directions for use and, in any case, should appear on the label when there is sufficient label space in addition to mandatory label information.

Subpart B—Warning and Caution Statements for Drugs

§ 369.20 Drugs; recommended warning and caution statements.

ACETANILID.

Warning—Do not exceed recommended dosage. Overdosage or continued use may result in serious blood disturbances.

ACETOPHENETIDIN CONTAINING PREPARATIONS. (See § 201.309 of this chapter.)

Warning—This medication may damage the kidneys when used in large amounts or for a long period of time. Do not take more than the recommended dosage, nor take regularly for longer than 10 days without consulting your physician.

ANESTHETICS FOR EXTERNAL USE (LOCAL ANESTHETICS). (See also § 310.201(a)(19) and (23) of this chapter.)

Caution—Do not use in the eyes. Not for prolonged use. If the condition for which this preparation is used persists or if a rash or irritation develops, discontinue use and consult physician.

ANTIHISTAMINICS FOR EXTERNAL USE (EXCEPT PREPARATIONS FOR OPHTHALMIC USE).

Caution—Do not use in the eyes. If the condition for which this preparation is used persists or if a rash or irritation develops, discontinue use and consult physician.

ANTI-HISTAMINICS, ORAL. (See also § 310.201(a)(4) and (a)(24) of this chapter.)

Caution—This preparation may cause drowsiness. Do not drive or operate machinery while taking this medication. Do not give to children under 6 years of age or exceed the recommended dosage unless directed by physician.

The reference to drowsiness is not required on preparations for the promotion of sleep or on preparations that are shown not to produce drowsiness.

ANTIPYRINE.

Warning—Do not exceed recommended dosage. If skin rash appears, discontinue use and consult physician. ANTISEPTICS FOR EXTERNAL USE.

Caution—In case of deep or puncture wounds or serious burns, consult physician. If redness, irritation, swelling, or pain persists or increases or if infection occurs discontinue use and consult physician.

The reference to wounds and burns is not required on preparations intended solely for diaper rash.

ARSENIC PREPARATIONS.

Warning—Frequent or prolonged use may cause serious injury. Do not exceed recommended dosage. Keep out of the reach of children.

BELLADONNA PREPARATIONS AND PREPARATIONS OF ITS ALKALOIDS (ATROPINE, HYOSCYAMINE, AND SCOPOLAMINE (HYOSCINE); HYOSCYAMUS, STRAMONIUM, THEIR DERIVATIVES, AND RELATED DRUG PREPARATIONS.

Warning—Not to be used by persons having glaucoma or excessive pressure within the eye, by elderly persons (where undiagnosed glaucoma or excessive pressure within the eye occurs most frequently), or by children under 6 years of age, unless directed by a physician. Discontinue use if blurring of vision, rapid pulse, or dizziness occurs. Do not exceed recommended dosage.

Not for frequent or prolonged use. If dryness of the mouth occurs, decrease dosage. If eye pain occurs, discontinue use and see your physician immediately as this may indicate undiagnosed glaucoma.

In the case of scopolamine or scopolamine aminoxide preparations indicated for insomnia, the portion of the above warning that reads “children under 6 years of age” should read instead “children under 12 years of age”. BORIC ACID (POWDERED, CRYSTALLINE, OR GRANULAR).

Warning—Do not use as a dusting powder, especially on infants, or take internally. Use only as a solution. Do not apply to badly broken or raw skin, or to large areas of the body.

BROMIDES.

Caution—Use only as directed. Do not give to children or use in the presence of kidney disease. If skin rash appears or if nervous symptoms persist, recur frequently, or are unusual, discontinue use and consult physician.

CARBOLIC ACID (PHENOL) PREPARATIONS (MORE THAN 0.5 PERCENT) FOR EXTERNAL USE.

Warning—Use according to directions. Do not apply to large areas of the body. If applied to fingers or toes, do not bandage.

CATHARTICS AND LAXATIVES—IRRITANTS AND OTHER PERISTALTIC STIMULANTS.

Warning—Do not use when abdominal pain, nausea, or vomiting are present. Frequent or prolonged use of this preparation may result in dependence on laxatives.

Mercury preparations should have added to the “frequent use” statement, the words “and serious mercury poisoning”.

Phenolphthalein preparations should bear, in addition to the general warning, the following statement:

Caution—If skin rash appears, do not use this or any other preparation containing phenolphthalein.

See also Mineral Oil Laxatives.

CHLORATES: MOUTH WASH OR GARGLE.

Avoid swallowing.

COBALT PREPARATIONS (See also §250.106 of this chapter.)

Warning—Do not exceed the recommended dosage. Do not administer to children under 12 years of age unless directed by physician. Do not use for more than 2 months unless directed by physician.

This warning is not required on articles containing not more than 0.5 milligram of cobalt as a cobalt salt per dosage unit and which recommend administration of not more than 0.5 milligram per dose and not more than 2 milligrams per 24-hour period.

“COUGH-DUE-TO-COLD” PREPARATIONS. (See also §310.201(a)(20) of this chapter.)

Warning—Persons with a high fever or persistent cough should not use this preparation unless directed by physician.

COUNTERIRRITANTS AND RUBEFACIENTS.

Caution—Do not apply to irritated skin or if excessive irritation develops. Avoid getting into the eyes or on mucous membranes.

If offered for use in arthritis or rheumatism, in juxtaposition therewith, the statement:

Caution—If pain persists for more than 10 days, or redness is present, or in conditions affecting children under 12 years of age consult a physician immediately.

See also “Salicylates” in this section for additional warnings for preparations containing methyl salicylate.

CREOSOTE, CRESOLS, GUAIACOL, AND SIMILAR SUBSTANCES IN PREPARATIONS FOR EXTERNAL USE.

Caution—Do not apply to large areas of the body.

CREOSOTE, CRESOLS, GUAIACOL, AND SIMILAR SUBSTANCES IN DOUCHE PREPARATIONS.

Warning—The use of solutions stronger than those recommended may result in severe local irritation, burns, or serious poisoning. Mix as directed before pouring into douche bag. Do not use more often than twice weekly unless directed by physician.

DENTURE RELINERS, PADS, AND CUSHIONS.

Warning—For temporary use only. Long-term use of this product may lead to faster bone loss, continuing irritation, sores, and tumors. For Use Only Until a Dentist Can Be Seen.

DENTURE REPAIR KITS.

Warning—For emergency repairs only. Long-term use of home-repaired dentures may cause faster bone loss, continuing irritation, sores, and tumors. This kit for emergency use only. See Dentist Without Delay.

DOUCHE PREPARATIONS.

Warning—Do not use more often than twice weekly unless directed by physician.

See also Creosote * * * Douche for additional warning.

DRESSINGS, PROTECTIVE SPRAY-ON TYPE. (See also §310.201(a) (11) and (18) of this chapter.)

Warning—In case of deep or puncture wounds or serious burns consult physician. If redness, irritation, swelling or pain persists or increases or if infection occurs consult physician. Keep away from eyes or other mucous membranes. Avoid inhaling.

See also Dispensers Pressurized by Gaseous Propellants * * * for additional warnings to be included for products under pressure.

IODINE AND IODIDES (ORAL).

Caution—If a skin rash appears, discontinue use and consult physician.

MERCURY PREPARATIONS FOR EXTERNAL USE.

Warning—Discontinue use if rash or irritation develops or if condition for which used persists. Frequent or prolonged use, or application to large areas may cause serious mercury poisoning.

MINERAL OIL LAXATIVES. (See also §201.302 of this chapter.)

Caution—Take only at bedtime. Avoid prolonged use. Do not administer to infants or young children, in pregnancy, or to bedridden or aged patients unless directed by physician.

NASAL PREPARATIONS: VASO-CONSTRICTORS (PHENYL-PROPANOLAMINE).

§ 369.20

21 CFR Ch. I (4-1-08 Edition)

Caution—Do not exceed recommended dosage.

NUX VOMICA AND STRYCHNINE PREPARATIONS.

“Do not use more than the recommended dosage. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.”

OPHTHALMIC PREPARATIONS. (See also §200.50 of this chapter.)

Boric acid offered for use in the preparation of ophthalmic solutions should bear the statement: Prepare solution by boiling in water. Store in a sterile container. Prepare sufficient for one day’s use and discard unused portion.

PHENACETIN-CONTAINING PREPARATION. (See acetophenetidin.)

PHENYLPROPANOLAMINE HYDROCHLORIDE PREPARATIONS, ORAL.

Caution—Individuals with high blood pressure, heart disease, diabetes, or thyroid disease should use only as directed by physician.

POTASSIUM PERMANGANATE AQUEOUS SOLUTIONS (CONTAINING NOT MORE THAN 0.04 PERCENT POTASSIUM PERMANGANATE). (See §250.108 of this chapter.)

Warning—For external use on the skin only. Severe injury may result from use internally or as a douche. Avoid contact with mucous membranes.

QUININE AND OTHER CINCHONA DERIVATIVES (EXCEPT FOR USE IN MALARIA).

Caution—Discontinue use if ringing in the ears, deafness, skin rash, or visual disturbances occur.

RESINS, OLEORESINS, AND VOLATILE OILS.

Caution—If nausea, vomiting, abdominal discomfort, diarrhea, or skin rash occurs, discontinue use and consult physician.

RESORCINOL (NOT THE MONOACETATE) HAIR PREPARATIONS.

Caution—Excessive use of this preparation may temporarily discolor blond, white, or red hair.

SALICYLATES, INCLUDING ASPIRIN AND SALICYLAMIDE (EXCEPT METHYL SALICYLATE, EFFERVESCENT SALICYLATE PREPARATIONS, AND PREPARATIONS OF AMINOSALICYLIC ACID AND ITS SALTS). (See also §201.314 of this chapter.)

“Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away;” or “Keep out of reach of children.”

If the article is an aspirin preparation, it should bear the first of the above two warning statements. In either case, the above information should appear on the label.

Caution—For children under 3 years of age, consult your physician; or

Caution—For younger children, consult your physician.

One of the two immediately preceding caution statements is required on the label of all aspirin tablets, but such a statement is not required on the labels of other salicylates clearly offered for administration to adults only.

If offered for use in arthritis or rheumatism, in juxtaposition therewith, the statement:

Caution—If pain persists for more than 10 days, or redness is present, or in conditions affecting children under 12 years of age, consult a physician immediately.

SALICYLATES: METHYL SALICYLATE (WINTERGREEN OIL). (See also §§201.303 and 201.314 of this chapter.)

“Do not use otherwise than as directed. Keep out of reach of children to avoid accidental poisoning. If swallowed, get medical help or contact a Poison Control Center right away.”

If the preparation is a counter-irritant or rubefacient the statement:

Caution—Discontinue use if excessive irritation of the skin develops. Avoid getting into the eyes or on mucous membranes.

If offered for use in arthritis or rheumatism, in juxtaposition therewith, the statement:

Caution—If pain persists for more than 10 days, or redness is present, or in conditions affecting children under 12 years of age consult a physician immediately.

SILVER.

Caution—Frequent or prolonged use of this preparation may result in permanent discoloration of skin and mucous membranes.

SODIUM PERBORATE MOUTHWASH AND GARGLE AND TOOTHPASTE.

Caution—Discontinue use if irritation or inflammation develops, or increases. Avoid swallowing.

SULFONAMIDE NOSE DROPS.

Caution—Do not use if a known allergy to sulfonamide drugs exists.

SULFUR PREPARATION FOR EXTERNAL USE.

Caution—If undue skin irritation develops or increases, discontinue use and consult physician.

THROAT PREPARATIONS FOR TEMPORARY RELIEF OF MINOR SORE THROAT: LOZENGES, TROCHES, WASHES, GARGLES, ETC. (See also §201.315 of this chapter.)

Warning—Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult physician promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by physician.

TOOTHACHE PREPARATIONS.

For temporary use only until a dentist can be consulted.

ZINC STEARATE DUSTING POWDERS.

“Keep out of reach of children; avoid inhaling. If swallowed, get medical help or contact a Poison Control Center right away.”

[39 FR 11745, Mar. 29, 1974, as amended at 40 FR 8917, Mar. 3, 1975; 40 FR 13496, Mar. 27, 1975; 41 FR 10885, Mar. 15, 1976; 51 FR 27760, Aug. 1, 1986; 51 FR 35340, Oct. 2, 1986; 52 FR 15893, Apr. 30, 1987; 52 FR 30057, Aug. 12, 1987; 52 FR 47324, Dec. 11, 1987; 53 FR 7093, Mar. 4, 1988; 55 FR 31783, Aug. 3, 1990; 57 FR 58376, Dec. 9, 1992; 59 FR 43412, Aug. 23, 1994; 64 FR 13296, Mar. 17, 1999; 68 FR 18882, April 17, 2003; 68 FR 34293, June 9, 2003]

§ 369.21 Drugs; warning and caution statements required by regulations.

ACETAMINOPHEN (N-ACETYL-*p*-AMINOPHENOL) (See §310.201(a)(1) of this chapter.)

Warning—Do not give to children under 3 years of age or use for more

than 10 days unless directed by a physician.

If offered for use in arthritis, or rheumatism, in juxtaposition therewith, the statement:

Caution—If pain persists for more than 10 days, or redness is present, or in conditions affecting children under 12 years of age consult a physician immediately.

ALCOHOL RUBBING COMPOUND. (See 26 CFR 182.855(a)(5); The National Formulary, Tenth Edition 1955, pp. 27-28; and section 502(g) of the act).

Warning—For external use only. If taken internally serious gastric disturbances will result.

ANTIHISTAMINICS, ORAL (PHENYLTOLOXAMINE DIHYDROGEN CITRATE AND CHLOROTHEN CITRATE PREPARATIONS). (See §310.201(a)(4) and (a)(24) of this chapter.)

Caution—This preparation may cause drowsiness. Do not drive or operate machinery while taking this medication. Do not give to children under 6 years of age or exceed the recommended dosage unless directed by physician.

If offered for symptoms of colds, the statement:

Caution—If relief does not occur within 3 days, discontinue use and consult physician.

DICYCLOMINE HYDROCHLORIDE WITH AN ANTACID. (See §310.201(a)(8) of this chapter.)

Warning—Do not exceed the recommended dosage. Do not administer to children under 12 years of age or use for a prolonged period unless directed by physician, since persistent or recurring symptoms may indicate a serious disease requiring medical attention.

DIPHEMANIL METHYLSULFATE FOR EXTERNAL USE. (See §310.201(a)(22) of this chapter.)

Caution—If redness, irritation, swelling, or pain persists or increases, discontinue use and consult physician.

DRUGS IN DISPENSERS PRESSURIZED BY GASEOUS PROPELLANTS. (See also §310.201(a)(11) and (18) of this chapter.)

The warnings herein shall appear prominently and conspicuously, but in