

Subpart H—Albumin (Human)**§ 640.80 Albumin (Human).**

(a) *Proper name and definition.* The proper name of the product shall be Albumin (Human). The product is defined as a sterile solution of the albumin derived from human plasma.

(b) *Source material.* The source material of Albumin (Human) shall be plasma recovered from Whole Blood prepared as prescribed in §§ 640.1 through 640.5, or Source Plasma prepared as prescribed in §§ 640.60 through 640.76.

(c) *Additives in source material.* Source material shall not contain an additive unless it is shown that the processing method yields a final product free of the additive to such extent that the continued safety, purity, potency, and effectiveness of the final product will not be adversely affected.

[42 FR 27582, May 31, 1977, as amended at 50 FR 4140, Jan. 29, 1985; 64 FR 26286, May 14, 1999]

§ 640.81 Processing.

(a) *Date of manufacture.* The date of manufacture shall be the date of final sterile filtration of a uniform pool of bulk solution.

(b) *Processing method.* The processing method shall not affect the integrity of the product, and shall have been shown to yield consistently a product which is safe for intravenous injection.

(c) *Microbial contamination.* All processing steps shall be conducted in a manner to minimize the risk of contamination from microorganisms, pyrogens, or other impurities. Preservatives to inhibit growth of microorganisms shall not be used during processing.

(d) *Storage of bulk fraction.* Bulk concentrate to be held more than 1 week prior to further processing shall be stored in clearly identified closed vessels at a temperature of -5°C or colder. Any other bulk form of the product, exclusive of the sterile bulk solution, to be held more than 1 week prior to further processing shall be stored in clearly identified closed vessels at a temperature of 5°C or colder. Any bulk fraction to be held one week or less prior to further processing shall be stored in clearly identified closed vessels at a temperature of 5°C or colder.

(e) *Heat treatment.* Heating of the final containers of Albumin (Human) shall begin within 24 hours after completion of filling. Heat treatment shall be conducted so that the solution is heated continuously for not less than 10, or more than 11 hours, at an attained temperature of $60\pm 0.5^{\circ}\text{C}$.

(f) *Stabilizer.* Either 0.08 ± 0.016 millimole sodium caprylate, or 0.08 ± 0.016 millimole sodium acetyltryptophanate and 0.08 ± 0.016 millimole sodium caprylate per gram of protein shall be present as a stabilizer(s). Calculations of the stabilizer concentration may employ the labeled value for the protein concentration of the product as referred to in § 640.84(d).

(g) *Incubation.* All final containers of Albumin (Human) shall be incubated at 20 to 35°C for at least 14 days following the heat treatment prescribed in paragraph (e) of this section. At the end of this incubation period, each final container shall be examined and all containers showing any indication of turbidity or microbial contamination shall not be issued. The contents of turbid final containers shall be examined microscopically and tested for sterility. If growth occurs, organisms shall be identified as to genus, and the material from such containers shall not be used for further manufacturing.

[42 FR 27582, May 31, 1977, as amended at 50 FR 4140, Jan. 29, 1985; 64 FR 26286, May 14, 1999; 65 FR 13679, Mar. 14, 2000; 65 FR 52018, Aug. 28, 2000]

§ 640.82 Tests on final product.

Tests shall be performed on the final product to determine that it meets the following standards:

(a) *Protein concentration.* Final product shall conform to one of the following concentrations: 4.0 ± 0.25 percent; 5.0 ± 0.30 percent; 20.0 ± 1.2 percent; and 25.0 ± 1.5 percent solution of protein.

(b) *Protein composition.* At least 96 percent of the total protein in the final product shall be albumin, as determined by a method that has been approved for each manufacturer by the Director, Center for Biologics Evaluation and Research, Food and Drug Administration.

(c) *pH.* The pH shall be 6.9 ± 0.5 when measured in a solution of the final